

Medicina Sportiva

Journal of the Romanian Sport Medicine Society

Vol. XI, No. 4 – 2015

15.12.2015

EDITORIAL STAFF

Assoc. Prof. Mirela Maria Vasilescu - *Editor-in-Chief*

Assoc. Prof. Anca Ionescu - *Scientific Editor*

EDITORIAL BOARD

Apostol Adela - University of Medicine and Pharmacy Carol Davila, Bucharest, Romania

Busneag Răzvan – National Institute of Sports Medicine, Bucharest, Romania

Fabian Klein – University of Medicine and Pharmacy Carol Davila, Bucharest, Romania

Georgescu Mariana - National Institute of Sports Medicine, Bucharest, Romania

Panait Gabriel – Sports Medicine Department, Vitan Policlinic, Bucharest, Romania

Popescu Alin – Rugby Romanian Federation, Bucharest, Romania

Siirghi Brandusa - University of Medicine and Pharmacy Carol Davila, Bucharest, Romania

SCIENTIFIC REVIEW BOARD

Avramescu Taina Elena - University of Craiova, Romania

Berteanu Mihai - University of Medicine and Pharmacy Bucharest, Rehabilitation Department, Romania

Bas Aslan Ummuhan – Pamukkale University, Denizli, Turkey

Branzaniuc Klara – University of Medicine, Tg Mures, Anatomy Department, Romania

Cavlak Ugur - Pamukkale University, School of Physical Therapy and Rehabilitation, Denizli, Turkey

Derevenco Petru - Romanian Academy of Medical Sciences, Romania

Dikic Nenad - Anti-Doping Agency of Serbia, Sports Academy, Beograd, Serbia

Dimitrova Diana – Sports Medicine Department, National Sports Academy, Sofia, Bulgaria

Drosescu Paula – University A.I.Cuza, Faculty of Physical Education and Sport, Iasi, Romania

Emin Ergen – University of Medicine, Ankara, EFSMA Executive Board, FIMS Executive Board, Turkey

Gamze Ekici - Ahi Evran University, School of Physical Therapy and Rehabilitation, Kirsehir, Turkey

Georgescu Luminita - University of Pitesti, Romania

Kürşad Serthbaş – Sakarya University, Physical Education and Sport Department, Turkey

Mazzeo Filomena - Parthenope University, Naples, Italy

Nastis Konstantinos - Aristotle University of Thessaloniki, Greece

Nestianu Valeriu - Romanian Academy of Medical Sciences, Romania

Nica Adriana – University of Medicine and Pharmacy, Bucharest, Romania

Oravitan Mihaela – West University of Timisoara, Romania

Rusu Ligia - University of Craiova, Romania

Domenico Tafuri - Parthenope University, Naples, Italy

ISSN 1841-0162

Publisher “Universitaria”, Craiova, Romania

Editorial Office Address: University of Craiova, Physical Education and Sports Faculty,
Brestei 156, 200177, Craiova, Romania

Site adress:

<http://www.medicinasportiva.ro>

Technical editor

Eng. Aurora Beldiman (University of Craiova)

GUIDE FOR AUTHORS

Medicina Sportiva, a publication of Romanian Sports Medicine Society, is a quarterly, peer reviewed journal which publishes original research, reviews and case reports on all areas of sports medicine and exercise science (exercise physiology, sports biomechanics, sports biochemistry, sports nutrition, diseases and exercise, sports injuries and sports traumatology, kinesiology and kinetotherapy).

The journal includes editorials, reviews, original papers, clinical studies, news in sport medicine and exercise science. Medicina Sportiva is currently indexed

Manuscript Preparation

All manuscripts must be submitted in English. Manuscripts are considered if they are submitted only to Medicina Sportiva and therefore should not be under consideration for publication elsewhere, either in part or in whole. Only the authors are responsible for the scientific context and legal aspects of the articles.

Reviews include theoretical and practical aspects that must to be report to actual researches in sport medicine. Size: the article must to have no more then 12 pages. Format: the text must to be compact and can has few subtitles.

Original papers include fundamentals research and clinical research regarding diagnostic and treatment. Size: 8-10 pages, 4-5 imagining documents (graphics, images, etc.). Format: the text must to be structure in introduction (present stage of the research), aim and scope of study, material and method, results, discussion, conclusions (must to present personal conclusions).

Case reports include specific clinical cases that provide relevant information on diagnosis and therapy of a particular case that proves unique to clinical sports medicine. Content: clinical aspects and researches regarding morphologic evaluation and clinic assessment, therapeutic conclusions and evolution. Size: 3-5pages, 1-2 original demonstrations and researches. Format: case report, discussion and comparative study in report with actualities studies.

Organization of the manuscript

Each manuscript should follow the sequence:

Title

Authors and affiliation

Abstract and Keywords

Main text (Introduction, Methods, Results, Discussion)

Acknowledgements,

References

Corresponding author

Authors' details

Text style of the manuscript

All manuscripts must to be edited in MS Word:

- page setup format A4, all margins at 2cm on all sides, single line spacing;

- font style Times New Roman, without capital letter,

- size:

14 for title (bold),

12 for authors (regular),

12 for affiliation (italic)

10 for abstract (regular),

10 for key words (italic),

11 for main text and (regular),

10 for references (regular, except the title of books and journals which will be written italic);

- paragraph:

indentation = 0,

alignment left for title, authors, affiliation, key words and justified for abstract, main text, acknowledgements and references;

- line spacing = single, column = one.

Tables, Figures and Schemes are to be typed on separate pages at the end of the manuscript. Ensure that each illustration or table has a caption attached, which should comprise a brief title and a description. Explain all symbols and abbreviations used. Units of measurement, symbols and abbreviations must conform to international standards. Do not imbed figures or tables into the text.

The tables, figures and schemes must be in the final published size, not oversized. Illustrations and tables that have appeared elsewhere must be accompanied by written permission to reproduce them from the original publishers. This is necessary even if you are an author of the borrowed material. Borrowed material should be acknowledged in the captions in the exact wording required by the copyright holder. If not specified, use this style: "Reproduced by kind permission of . . . (publishers) from . . . (reference)". Identifiable clinical photographs must be accompanied by written permission from the patient. The first time an abbreviation appears in the text, it should be preceded by the words for which it stands.

Title: should be simple, concise and informative. Capitalize the first word only.

Author names should appear as used for conventional publication, with first and middle names or initials followed by surname. If multiple authors have the same affiliation, one listing of the affiliation should be used, preceded by the full list of those authors on the line above.

Abstract: begin the section with the word **Abstract** in bold font. The authors prepare a structured abstract of not more than 300 words. The abstract should be an explicit summary of the paper and must include (at least) four headings: aim, methods, results, conclusion.

Key words. Immediately after the abstract, provide a maximum of 3–6 keywords. Key words should refer to the terms from Medical Subject Headings (MeSH) of MEDLINE/PubMed.

Acknowledgments: Acknowledgments should be included before the References section.

References: in the text identify references by Arabic numeral in brackets, using Oxford system. The list of references should include only those publications which are cited in the text. References should be presented in consecutive order as they are cited in the text. References should be cited in the text by placing sequential numbers in brackets (for example, (1), (3-5)). The four six authors should be presented. No more than 30 references will be accepted.

Examples:

a) Journals: name of authors, initial of surname, years of publish in brackets, title of paper, title of periodical (italic), volume, page (Ryder, JW, Bassel DUBY R, Olson EN, Zierath, JR (2003). Skeletal muscle reprogramming by activation

of calcineurin improves insulin action on metabolic pathways. *J. Biol. Chem.* 278, 298 304);

b) Books: name of authors, years of publish in brackets, title of papers or capitols, title of book or monograph (italic), editors, city, pages (Booth FW, Baldwin KM (1995). Muscle plasticity: energy demanding and supply processes. In: *Handbook of physiology*. Rowell LB, Shepherd JT (eds). Oxford University Press, New York, pp 1076-1121).

c) Proceedings: name of authors, initial of surname, years of publish in brackets, title of paper, title of periodical (italic), editors, city, pages.

Corresponding author should contain all information necessary for an effective correspondence (postal addresses, e-mail, fax and telephone number).

Authors' details

It must include a brief biography, approximately 70 words per author, for each author including academic title/degrees, institution affiliation, research focus, e-mail address.

Ethical standards

Research carried out on humans must be in compliance with the Helsinki Declaration, adopted by the 18th World Medical Assembly, Helsinki, Finland, June 1964 and any experimental research on animals must follow internationally recognized guidelines.

By submitting a manuscript to our journal, each author explicitly confirms that the manuscript meets the highest ethical standards for authors and coauthors. The manuscript should contain a statement that has been approved by the appropriate ethical committees related to the institution(s) in which it was performed and that subjects gave informed consent to the work.

Studies involving experiments with animals must state that their care was in accordance with institution guidelines. Patients' and volunteers' names, initials, and hospital numbers should not be used. In a case report, the subject's written consent should be provided. It is the author's responsibility to ensure all appropriate consents have been obtained. Patients have a right to privacy. Therefore identifying information, including patients images, names, initials, or hospital numbers, should not be included in videos, recordings, written descriptions, photographs, and pedigrees unless the information is essential for scientific purposes and you have obtained written informed consent for publication in print and electronic form from the patient (or parent, guardian or next of kin where applicable). If such consent is made subject to any conditions,

Universitaria Publisher must be made aware of all such conditions. Written consents must be provided to Universitaria Publisher on request. Even where consent has been given, identifying details should be omitted if they are not essential. If identifying characteristics are altered to protect anonymity, such as in genetic pedigrees, authors should provide assurance that alterations do not distort scientific meaning and editors should so note. If such consent has not been obtained, personal details of patients included in any part of the paper and in any supplementary materials (including all illustrations and videos) must be removed before submission.

Copyright agreement and plagiarism

Manuscripts are welcomed from all countries, and may be submitted for consideration on the understanding that they have not been previously published elsewhere, either in print or electronically, and that they are not under consideration by any other publisher.

Authors wishing to include figures or text passages that have already been published elsewhere are required to obtain permission from the copyright holder(s) and to include evidence that such permission has been granted when submitting their papers. Authors are required to ensure that no material submitted as part of a manuscript infringes existing copyrights, or the rights of a third party.

The author who submits the manuscript for publication should ensure that all coauthors are included on the paper, and that all coauthors have seen the final version of the article and have agreed to its submission for publication. Only, the authors are responsible for the scientific context and legal aspects of publishing the manuscript.

Any material received without such evidence will be assumed to originate from the authors. The authors are responsible for obtaining permission from copyright holders for reproducing any illustrations, tables, figures or lengthy quotations previously published elsewhere. Articles published in *Medicina Sportiva* may not be published elsewhere without written permission from the publisher.

If evidence of plagiarism is found before or after acceptance or after publication of the paper, the author will be offered a chance to defend his/her paper. If the arguments are not found to be satisfactory, the manuscript will be retracted and Authors found to have been guilty of plagiarism will no longer have papers accepted for publication in *Journal*.

The Editorial and Peer Review Process

All papers submitted to *Medicina Sportiva* are subject to peer review. Manuscripts must not be under simultaneous consideration by any other publication, before or during the peer-review process. Upon receipt of a manuscript the editor will notify the corresponding author that the submission has been received.

Received manuscripts are first examined by the *Medicina Sportiva* editors. Manuscripts with insufficient priority for publication are rejected. Manuscripts not prepared in the advised style will be sent back to authors without scientific review.

The manuscript will undergo a blinded review process. Under this process the editor will forward (electronically) the manuscript to two independent reviewers, members of the Scientific Board.

The editors encourage authors to suggest the names of potential reviewers, but they reserve the right of final selection. Normally the review process will take approximately 8 weeks.

After the review process has been completed the main author will be notified via e-mail either of acceptance, revision or rejection of the manuscript. If revision was recommended, this should be completed and returned to the editor within 2 weeks.

Editors may make necessary changes to articles in accordance with guidelines for authors.

Authors' rights

As an author you (or your employer or institution) may do the following:

- make copies (electronic) of the article for your own personal use, including for your own classroom teaching use
- present the article at a meeting or conference and to distribute copies of the article to the delegates attending such a meeting
- for your employer, if the article is a “work for hire”, made within the scope of your employment, your employer may use all or part of the information in the article for other intra-company use (e.g., training)
- retain patent and trademark rights and rights to any processes or procedure described in the article
- include the article in full or in part in a thesis or dissertation (provided that this is not to be published commercially)
- use the article or any part thereof in a printed compilation of your works, such as collected writings or lecture notes (subsequent to publication of your article in the journal)

•prepare other derivative works, to extend the article into book-length form, or to otherwise re-use portions or excerpts in other works, with full acknowledgement of its original publication in the journal.

Submission of manuscripts

Authors are encouraged to submit their manuscripts via e-mail to editors, by e-mail *medsprt@yahoo.com*

Copyright Transfer Statement

Article title	
Article authors	
Full name of the corresponding author	
Title	
Affiliation	
Address	
City, country	
Telephone	
E-mail	

The copyright to this article is transferred to MEDICINA SPORTIVA (ISSN 1841-0162) if and when the article is accepted for publication.

The undersigned hereby transfers any and all rights in and to the paper including without limitation all copyrights to MEDICINA SPORTIVA.

The authors of this article declare that:

1. are responsible for the scientific context and legal aspects of the articles.
2. this manuscript has not been published in the same form elsewhere.
3. it will not be submitted anywhere else for publication before or during the peer-review process, acceptance or rejection by the journal MEDICINA SPORTIVA.
4. are responsible for obtaining permission from copyright holders for reproducing any illustrations, tables, figures or lengthy quotations previously published elsewhere.
5. the article published in Medicina Sportiva may not be published elsewhere without written permission from the publisher.
6. transfer the unlimited rights of publication of the above mentioned paper in whole to the journal MEDICINA SPORTIVA.
7. the copyright transfer covers the exclusive right to reproduce and distribute the article in all the formats

The corresponding author signs for and accepts responsibility for releasing this material on behalf of any and all co-authors.

After submission of this agreement signed by the corresponding author, any changes of the authors list will not be accepted.

Date:

Corresponding author:

Posttraumatic peripheral neuropathic pain and CRPS in the upper limb

Constantinovic Mariana Isabela¹, Mologhianu Gilda¹, Petru Armean¹, Roxana Nartea², Mirela Vasilescu³, Nica Sarah Adriana¹

¹Carol Davila University of Medicine and Pharmacy, Bucharest, Romania,

²Buzau County Hospital,

³Craiova University, Kinetotherapy and Sports Medicine Department, Craiova, Romania

Abstract. The term of "complex regional pain syndromes" (CRPS) was introduced by IASP (*International Association for the Study of Pain*) in 1994 and represents a group of algic disorders that may develop as a disproportional consequence of a minor trauma (contusion, fracture, postsurgical), usually in one extremity. There have been described 2 types of CRPS I, previously known as sympathetic dystrophy, develops following trauma with minor lesions and without obvious neurological impairment (with fractures, sprains, contusions or cutaneous lesions). CRPS II or causalgia has similar signs and symptoms and develops after trauma with important peripheral nerve injuries (10-12). Neuropathic pain therapy is often inefficient regarding nociceptive pain. Knowledge of the patient's medical history related to neuropathic pain and using validated assessment tools are essential for differentiating neuropathic pain from nociceptive pain and to estimate the importance of the neuropathic component of pain from various pain syndromes.

In 2007, International Society for The Study of Pain (IASP) proposed the following definition for neuropathic pain as "pain caused by a lesion or disease of the somatosensory system" (13). "Diseases" are pathological processes such as inflammatory, autoimmune or dysfunction of ion channels. "Lesions" are minor detectable lesions. Diseases and lesions of nervous system elements others than somatosensory component determine nociceptive pain. Spasticity and muscle pain related to motor system diseases and lesions are not being associated with neuropathic pain.

Keywords: *nerve injuries, algic disorders, sympathetic dystrophy.*

Neuropathic pain definition

Modern research on neuropathic pain showed that traumatic peripheral nerve injuries determine important changes of the sensory component of the nervous system. Those changes differentiate neuropathic pain from other chronic pain types, such as chronic nociceptive pain from osteoarthritis, in which case the nociception is not impaired (1).

Posttraumatic peripheral neuropathic pain results from mechanical trauma to the peripheral nerve most frequently related to car accidents, work accidents or in domestic context (2, 3). According to WHO, 3.200.000 experience posttraumatic disabilities after car accidents (4).

Besides classical neuropathic syndromes such as phantom limb pain, there are posttraumatic algic states with common clinical features. These syndromes previously known as reflex sympathetic dystrophy, Sudeck dystrophy, causalgia, shoulder-hand syndrome or algodystrophy are in present grouped under the umbrella term of "complex regional pain syndromes" (CRPS) (1). The term CRPS was introduced by IASP (*International Association for the Study of Pain*) in 1994 and represents a group of algic disorders that may develop as a disproportional consequence of a minor trauma (contusion, fracture, postsurgical), usually in one extremity (5-9).

There have been described 2 types of CRPS I, previously known as sympathetic dystrophy, develops following trauma with minor lesions and without obvious neurological impairment (with fractures, sprains, contusions or cutaneous lesions). CRPS II or causalgia has similar signs and symptoms and develops after trauma with important peripheral nerve injuries (10-12).

Neuropathic pain therapy is often inefficient regarding nociceptive pain. Knowledge of the patient's medical history related to neuropathic pain and using validated assessment tools are essential for differentiating neuropathic pain from nociceptive pain and to estimate the importance of the neuropathic component of pain from various pain syndromes (1).

In 2007, International Society for The Study of Pain (IASP) proposed the following definition of neuropathic pain as “pain caused by a lesion or disease of the somatosensory system” (13). “Diseases” are pathological processes such as inflammatory, autoimmune or dysfunction of ion channels. “Lesions” are minor detectable lesions.

Diseases and lesions of nervous system elements others than somatic-sensory component determine nociceptive pain. Spasticity and muscle pain related to motor system diseases and lesions are not being associated with neuropathic pain.

Also, is important to establish the peripheral or central origin of pain, taking into account the topography, clinical signs and pathophysiological mechanisms being different.

Classification

In clinical practice, neuropathic pain is classified according to the etiology and anatomical topography of the injury (12). There are 4 major types of nervous system impairment: peripheral painful neuropathies (for example, traumatic, ischemic, inflammatory, toxic, metabolic or hereditary), central type pain syndromes (for example, in stroke, multiple sclerosis, spinal injury), neuropathic complex pain disorders (CRPS) and mixed pain syndromes (nociceptive and neuropathic pain). In many chronic pain syndromes there are involved both nociceptive and neuropathic processes. Those mechanisms can explain the various symptomatology (mixed pain syndromes) (1). The outcome can vary from relatively mild to chronic pathology, with a major impact on daily functioning and quality of life.

Clinical signs in complex regional pain syndromes

CRPS develops usually in the extremities, following fractures, surgery, prolonged immobilization, peripheral nerve injury, etc. It usually manifest in the distal part of the extremity, but pain may radiate proximally or to other body parts. Upper extremities are most frequently impaired. Pain is continuous, spontaneous, regional, like burning, and does not respect the anatomical distribution of a peripheral nerve (11). Symptoms can install gradually after the trauma (14-17). Immediately after the trauma, CRPS is difficult to detect because of the similar posttraumatic symptoms (18).

There was estimated that 50% of the patients with CRPS develop a chronic pain syndrome and disability (12). Pain is disproportional to the trauma severity and associates signs of neurogenic inflammation, osteoporosis and other signs such as (19-22): sensory disorders/dysfunction (burning pain, mechanical hyperalgesia, hyperesthesia, allodynia, sudomotor disorders (oedema and excessive perspiration) and vasomotor disorders (skin temperature and color changes); trophic changes of skin, hair and nails; motor function impairment: voluntary movement impairment, reduced range of motion, hypotonia, tremor, joint stiffness, coordination disorders, dystonia and muscle spasms.

Pain related to peripheral nerve injury has specific clinical features (23). If the injured nerve is a mixed peripheral nerve with a cutaneous branch there is an area of abnormal sensation. The maximum pain experienced by the patient corresponds to an area of sensory deficit, specific for neuropathic pain. Sensory deficit is usually related to nociceptive or thermal stimuli (1).

Paresthesia determine discomfort, but are not painful. Positive signs are sensations of permanent and intermittent spontaneous pain (are not induced by stimuli). Many patients with neuropathic pain reported also stimuli induced pain and hypersensitivity. Patients report frequently mechanical hypersensitivity and consequently, hypersensibility (1). There are two types of hypersensibility: allodynia - pain as a response to non-nociceptive stimuli, and hyperalgesia - painful hypersensitivity to nociceptive stimuli.

A small number of patients with peripheral nerve injury have a hypersensitive syndrome, without evident signs of sensory deficit. Usually, neuropathic pain is intermittent, like burning, with unusual paresthesia. Although these characteristics are not universally present in all neuropathic pain states, when they are, the diagnosis of neuropathic pain is probable (1).

Peripheral sensitization of primary afferents

Peripheral sensitization of primary afferent neurons has been described in patients with painful peripheral nerve injury. An example is decreased thermal algic perception. Cutaneous cold hypersensibility (hyperalgesia) is frequently in patients with posttraumatic neuralgia and in chronic complex regional pain syndromes (1).

Neuropathic pain persists more than nociceptive pain, is provoked by stimuli that are normally not painful and is influenced only by specific medication (14).

Neuropathic pain is severe, impairing patient's life quality, being associated with anxiety and depression, sleep disorders and disability (24, 14). It may develop after trauma with peripheral nerve injuries or after surgery (25, 3, 14). Even in patients with minor changes in neurologic tests, in peripheral nerve injuries neuropathic pain is spontaneous and determine areas of hypoaesthesia and/or hyperaesthesia (14).

Ciaramitaro et al. have published in 2010 an epidemiological study regarding traumatic peripheral nerve lesions and neuropathic pain. In the first year they analyzed 211 cases, most of them young males, the lesions being predominantly in the upper limb. 50% of them presented neuropathic pain, mild to severe in 79% of the patients. Most frequently pain was located in radial and ulnar nerve areas. The presence of neuropathic pain has been correlated with decreased quality of life in patients with posttraumatic neuropathies (3, 14).

Taylor et al. (2010) compared two groups of patients with postsurgical peripheral nerve lesions (median and cubital). Patients with neuropathic pain had severe sensory-motor deficit and impaired nerve recovery (26, 14).

Incidence and etiology of CRPS

The incidence of CRPS was analyzed in retrospective and prospective studies. They have been reported most frequently after distal radius fractures (27-29, 21, 19). Sandroni et al. reported an incidence rate of CRPS of 5.46/100.000 person years in the general population (Olmsted County, USA) (30, 8).

CRPS impairs 3 times more women than men and may develop in any age, with a higher incidence between ages of 50 and 70 years. Over 50% are posttraumatic. In 44% of cases, the trigger was a fracture (8). Most of the studies were prospective. The incidence was 1-2% in retrospective studies regarding distal radius fractures and up to 38% in prospective studies (31), probably due to a lack of uniform and specific diagnostic criteria (11). Other reported causes are sprains (18%) and surgery (12%) (8). Also, prolonged immobilization, peripheral nerve injury and compressive syndromes may determine CRPS (28).

Clinical signs, especially those of autonomic dysfunction and Leriche findings on pain relief through sympathectomy sustain the role of the nervous sympathetic system in CRPS etiology (32, 9).

Pain from complex regional pain syndromes has a component of neuropathic pain (33, 9), being generated and maintained by peripheral nerve sensory fibers and also is inflammatory, nociceptive and functional (9). Quantitative sensory testing suggests the role of inflammation in acute CRPS. Nociceptive pain may be caused by tissue lesions, as a result of the initial trauma or by secondary tissue lesions determined by edema and trophic changes, hypoxia and acidosis (34-35, 9).

CRPS pathophysiology

It has been suggested that trauma and rare diseases may determine disorders of the central nervous system. CRPS may be a disorder of sensory, autonomic and motor functions (11). Besides peripheral factors, CRPS may be also determined by neurogenic inflammation, immune factors and genetic predisposition (36-39, 11). Spatial distribution of pain, allodynia to mechanical and thermal stimuli and hyperalgesia suggests the role of both peripheral nervous system and central nervous system in the pathophysiology of complex regional pain syndromes (40, 41, 20, 9).

Some symptoms suggests that neurological dysfunction is not limited to the peripheral and autonomic nervous system, but there is also a distortion of body image and cortical function reorganization (42, 22), at the cortical level.

In the study of Pleger et al., cortical representation of painful hand was significantly reduced compared to contralateral normal hand (43, 11). Maihofner et al. reported that cortical representation of the normal contralateral hand was significantly more reduced than in the CRPS impaired hand, considering that is due to the reorganization of the primary somatosensory cortex (44, 11). It has been found that cortical reorganization is CRPS nonspecific, being reported also in other algic states (11) and it becomes normal along with the recovery (45, 11).

Therapeutical management of CRPS

The mainly therapeutic goals in CRPS are pain control and functional restoration. Findings of different studies showed that early diagnosis of CRPS and prompt initiation of individualized therapy are essential to improve patient outcomes. Because of the complex symptomatology, in the therapeutically management of CRPS there may be involved different specialists - in anesthesiology, surgery, neurology, rheumatology, physical medicine and rehabilitation (8).

Besides analgesic and antiinflammatory pharmacological therapy there may also be useful physical therapy, occupational therapy and psychotherapy (11). Active physiotherapy is essential for functional restoration (46, 11). Immobilization and splinting are the first physical modalities recommended. Also elevation of the distal part of the extremity, therapeutic massage and isometric exercises are gradually introduced into the therapeutic programme considering therapy must not exacerbate the pain (47). Hydrotherapy (alternating heat and cold baths), transcutaneous electrical nerve stimulation (TENS), and a stress loading program of traction and compression exercises have also been shown to be effective in CRPS (48, 47).

Functional neuroimaging studies reported the increase of motor cortical representation of the affected extremity (43) and reduced cortical representation of primary sensory cortex (44). It has been considered that the reorganization is reversible and that it is correlated with pain. Also, pain relief was correlated with a normalization of primary cortical sensory area (45).

Two interventions were developed based on the concept of sensory-motor information manipulation in the impaired extremity: mirror therapy (first introduced for phantom limb pain therapy) and prism adaptation (49- 53, 11, 22). Mirror therapy may help for magnifying range of motion and pain relief in patients with CRPS (50). Patients execute synchronal bimanual movements while they are looking the reflection of nonimpaired extremity in a mirror placed in the sagittal midline, having the illusion of a normal functional hand (50, 22).

It has been observed that pain is triggered by seeing an object approaching the impaired extremity (54, 22, 41) and may be increased when the patient looks at the object through magnifying lenses. Also it decrease when watching through lenses that reduce the images (55, 22). These findings suggested the pathological cortical reorganization and the existence of an extension of neural representation of peripersonal space (22).

Acknowledgements. This publication benefited from the financial support of the project Programme of Excellency in the Multidisciplinary Doctoral and Postdoctoral Research of Chronic Diseases, contract no. POSDRU/159/1.5/S/ 133377, beneficiary Gr. T. Popa University of Medicine and Pharmacy, Romania, Iasi, co-financed from the European Social Fund through the Sectoral Operational Programme Human Resources Development (SOP HRD) 2007-2013”.

References

1. Baron R, Neuropathic Pain (2009). A Clinical Perspective. *Handb Exp Pharmacol*; (194): 3-30. doi: 10.1007/978-3-540-79090-7_1.
2. Kouyoumdjian JA (2006). Peripheral nerve injuries: a retrospective survey of 456 cases. *Muscle and Nerve*; 34: 785–787.
3. Ciaramitaro P, Mondelli M, Logullo F, et al (2010). Traumatic peripheral nerve injuries: epidemiological findings, neuropathic pain and quality of life in 158 patients. *J Periph Nerv Sys*;15: 120–127.
4. Gin-Shaw SL, Jordan RC (2002). Multiple trauma. In: *Rosen's EmergencyMedicine: Concepts and Clinical Practice*. Marx R (Ed). New York, Mosby, pp 242–254.
5. Janig W, Baron R (2006). Is CRPS I a neuropathic pain syndrome? *Pain*; 120(3): 227-229.
6. Baron R, Janig W (2004). Complex regional pain syndromes – how do we escape the diagnostic trap? *Lancet*; 364: 1739–1741.
7. Merritt WH (2005). The challenge to manage reflex sympathetic dystrophy/complex regional pain syndrome. *Clin Plast Surg*; 32: 575–604.
8. de Mos M, De Bruijn AG, Huygen FJ, Dieleman JP, Stricker BHCh, Sturkenboom MCJM (2007). The incidence of complex regional pain syndrome: a population-based study. *Pain*; 129: 12–20.
9. Groeneweg G, Huygen FJPM, Coderre TJ, Zijlstra F (2009). Regulation of peripheral blood flow in Complex Regional Pain Syndrome: clinical implication for symptomatic relief and pain management. *BMC Musculoskeletal Disorders*; 10:116 doi:10.1186/1471-2474-10-116.
10. Stanton-Hicks M, Jänig W, Hassenbusch S, Haddock JD, Boas RA, Wilson PR (1995). Reflex sympathetic dystrophy: changing concepts and taxonomy. *Pain*; 1995;63:127–133.
11. Pelto-Vasenius Kia (2009). Posttraumatic Complex Regional Pain Syndrome (CRPS). *Suomen Ortopedia ja Traumatologia*; 32 (2):118-123.
12. Jensen TS, Baron R (2003). Translation of symptoms and signs into mechanisms in neuropathic pain. *Pain*; 102:1–8.
13. Treede R, Jensen T, Campbell JN, et al (2008). Neuropathic pain. Redefinition and a grading system for clinical and research purposes. *Neurology*; 70: 1630–1635.
14. Treede R, Jensen T, Campbell JN, et al (2008). Neuropathic pain. Redefinition and a grading system for clinical and research purposes. *Neurology*; 70: 1630–1635.