

The journal *Forum geografic. Studii și cercetări de geografie și protecția mediului* was founded in 2002, and it seeks to publish high quality research in the domains of geography, environment protection and other related domains. It addresses a range of issues, such as geomorphology, pedology, climatology, hydrology, human geography and environment. Its content is directed to a broad audience, including both academics and policymakers. The

papers selected for publication in the journal are subject to a review process using references from universities worldwide.

The journal is currently indexed by the following databases: **DOAJ** – Directory of Open Access Journals, **EBSCO** Publishing, **Index Copernicus** International, **Scipio** Scientific Publishing & Information Online, **CrossRef**, **ULRICHSWEB** - Global Serials Directory, **IGU** Journals Database.

Associate Editors:

Péter BAJMÓCY, Department of Economic and Human Geography, University of Szeged, H-6722 Szeged, Egyetem u. 2. Hungary

Slavoljub DRAGIĆEVIĆ, Faculty of Geography, Belgrade University, Studentski trg 3/3, Belgrade, Serbia

Vesna LUKIĆ, Demographic Research Centre, Institute of Social Sciences, Kraljice Natalije 45, Belgrade, Serbia

Nina NIKOLOVA, Faculty of Geology and Geography, "St. Kliment Ohridsky" University of Sofia, Tzar Osvoboditel Blvd. 15, Sofia, Bulgaria

Editorial Advisory Board:

Lucian BADEA, *The Institute of Geography, The Romanian Academy*

Dan BĂLTEANU, *The Institute of Geography, The Romanian Academy*

Zeljko BJELJAC, *Geographical Institute Jovan Cvijić, Serbia*

Sandu BOENGIU, *University of Craiova, Romania*

Léon BRENIG, *University of Brussels, Belgium*

Pompei COCEAN, *Babeş-Bolyai University, Romania*

Lóczy DÉNES, *University of Pécs, Hungary*

George ERDELI, *University of Bucharest, Romania*

Robert FOVELL, *University of California, USA*

Nelly HRISTOVA, *St. Kliment Ohridsky University of Sofia, Bulgaria*

Ioan IANOȘ, *University of Bucharest, Romania*

Mirela MAZILU, *University of Craiova, Romania*

Zvi Yehoshua OFFER, *Ben-Gurion University, Israel*

Maria PĂTROESCU, *University of Bucharest, Romania*

Maria RĂDOANE, *Ștefan cel Mare University, Romania*

Milan RADOVANOVIĆ, *Geographical Institute Jovan Cvijić, Serbia*

Recep EFE, *Balikesir University, Turkey*

Maria REDEY, *Eötvös Loránd University, Hungary*

Magdy TORAB, *Alexandria University, Egypt*

Alina VLADUȚ, *University of Craiova, Romania*

Nenad ŽIVKOVIĆ, *Belgrade University, Serbia*

Zbigniew ZWOLIŃSKI, *Adam Mickiewicz University (Poznan), Poland*

Editor-in-chief: **Sandu BOENGIU**, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Executive editor: **Liliana POPESCU**, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Assistant Editors: **Cristiana VÎLCEA**, **Oana IONUȘ**, **Amalia BĂDIȚĂ**, **Daniel SIMULESCU**

University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Cover photo: *the Râu-Șes erosion platform, Rânca - Parâng Mountains (photo Sandu Boengiu)*

For instructions for authors, subscription and all other information please visit our website

<http://forumgeografic.ro>

before submitting any papers please select the section *Publishing rules* from the About page and read thoroughly the submission instructions for authors

ISSN 1583-1523 (print)

ISSN 2067-4635 (online)

DOI prefix: 10.5775

Instructions for Authors

Article submission

In order to disseminate the research results in the field, researchers, scholars and professionals are welcome to submit an electronic version of the manuscript (in Microsoft Office Word format) to the editorial office (forum.geografic@gmail.com).

Submission requirements: The submission of an article for publication in our journal implies that the article has not been published before, nor it is being considered for publication in other journals. Authors are responsible for the content and the originality of their contributions. In order to be published, articles must be thoroughly researched and referenced.

IMPORTANT: All papers must be submitted in electronic format, only in English language.

Copyright statement

By submitting a scientific work to *Forum geografic* the submitters agree to declare the following:

- the submitted work belongs exclusively to the declared authors;
- the submitted work represents original scientific research;
- the submitted work has not been published or submitted for publishing to another journal;
- if the submitted work is published or selected for publishing in *Forum geografic*, the authors waive any patrimonial claims derived from their authorship for the submitted work; the authors retain the moral rights for their submitted work, as granted under the Romanian applicable law; also, the authors agree to refrain from ulterior submitting of the work to other journals.

The submitters agree to be solely held accountable in case of breaching the above terms and to defend the representatives of *Forum geografic* in the event of a lawsuit related to the submitted work.

When submitting a paper the authors are required to print, fill and send a scanned copy of this declaration.

Privacy statement

The submitted personal data, such as names or email addresses, are used only for the declared purpose of the *Forum geografic* journal (publishing original scientific research) and are not available to third parties.

Manuscripts are received at all times. However, in order to have your article published in the current year, the manuscripts must be submitted until the 15th of February for the first issue of the current year and until the 1st of September for the second issue.

Article format

All manuscripts must be edited entirely in English. Articles must include:

- Title
- Author's name(s). For each author you must mention the author's scientific title, his affiliation (institution) and e-mail address;

- Abstract (maximum 300 words);
- Keywords (not more than 5-6 words);
- Acknowledgments (if any);
- Main body of text (structured according to Introduction, Data & Methods, Results & Discussions, Conclusions);
- Illustrations (graphs, diagrams, maps, photos – should have indications of their positions in the text and title written in English) must be also submitted in electronic format, preferably in JPG, PNG or BMP format and must be referred to as Figures, which should be numbered with Arabic numbers.
- Tables must be numbered with Arabic numbers and should not repeat data available elsewhere in the text.
- References must be indicated in the text, between brackets and they must include the author's name and the date of the publication (Popescu, 2000). When three or more authors are referred, they will appear in the text as follows: (Popescu et al., 1997). References must be listed in alphabetical order at the end of the text.

The following style sheet is recommended:

- for journals:

Miletić, R., Lukić, V., & Miljanović, D. (2011). Deindustrialization and structural changes in commuting flows in Serbia. *Forum geografic*, X(2), 244-254. doi:10.5775/fg.2067-4635.2011.009.d

- for books:

Bran, F., Marin, D., & Simion, T. (1997). *Turismul rural. Modelul european*, Editura Economică, București

- for papers from conference proceedings:

Deci, E. L., Ryan, R. M., (1991), A motivational approach to self: Integration in personality. In R. Dienstbier (Ed.), *Nebraska Symposium on Motivation: Vol. 38. Perspectives on motivations* (pp. 237-288). Lincoln: University of Nebraska Press.

Review process

All the manuscripts received by the editors undergo an anonymous peer review process, necessary for assessing the quality of scientific information, the relevance to the field, the appropriateness of scientific writing style, the compliance with the style and technical requirements of our journal, etc. The referees are selected from the national and international members of the editorial and scientific board, as well as from other scholarly or professional experts in the field. The referees assess the article drafts, commenting and making recommendations. This process leads either to acceptance, recommendation for revision, or rejection of the assessed article. Editors reserve the right to make minor editorial changes to the submitted articles, including changes to grammar, punctuation and spelling, as well as article format, but no major alterations will be carried out without the author's approval. Before being published, the author is sent the proof of the manuscript adjusted by editors. If major revisions are necessary, articles are returned to the author so that he should make the proper changes. Authors are notified by email about the status of the submitted.

The Wind Regime of Romania – Characteristics, Trends and North Atlantic Oscillation Influences

Alfred VESPREMEANU-STROE^{1,*}, Sorin CHEVAL², Florin TĂTUI¹

¹ Faculty of Geography, University of Bucharest, N. Bălcesu Blvd, no 1, Bucharest, Romania

² National Administration of Meteorology, Bucharest-Ploiesti Road, no 97, 71552 Bucharest, Romania

* Corresponding author, fredi@geo.unibuc.ro

Received on <21-08-2012>, reviewed on <25-11-2012>, accepted on <08-12-2012>

Abstract

The present study attempts to develop a comprehensive perspective of the wind regime on the Romanian territory, focusing on the characteristics and tendencies encountered over the past 50 years, as well as on the NAO projection on it, using several data categories gathered from 167 meteorological stations. Based on the recorded multiannual averages and on the strong correlation ($r = 0.87$) established between altitude and wind speed in wind exposed areas, we created the map of mean (multi)annual wind speed. The highest aeolian potential corresponds to the Carpathians (7-10 m/s on the ridges and 3-7 m/s on the slopes and within valleys) and the Black Sea Shore (5-7 m/s); nevertheless, the two areas develop extreme values of the wind turbulence – maximum in the Carpathians and minimum in the coastal units. Tablelands in Moldova and Dobrogea, as well as the northern part of Bărăgan show ideal conditions for the development of aeolian parks (moderate and strong winds, low turbulences). On a country-size scale, the month with the highest mean wind speed is March, whereas August is the calmest one. For the first time, the map of resultant wind direction (DRV) was designed, expressing both the resultant wind orientation and its intensity (high intensity = low directional variability). Concerning NAO, a negative correlation is observed between its indices and the wind speed (mean speed, frequency and intensity of the stromic events) at a multiannual and multidecadal scale. The correlation coefficients present high values in the extra-carpathian areas and small or moderate values in the intra-carpathian areas. The positive (negative) NAO associated with low (high) cyclogenesis over the Mediterranean area induce low (high) winds over the Romanian territory due to diminished (enhanced) frequency of cyclones reaching Romanian regions.

Keywords: *wind regime, North Atlantic influences, wind direction*

Rezumat. Regimul vânturilor din România – Caracteristici, tendințe și influențele oscilațiilor nord-atlantice

Prezentul studiu încearcă să compună o perspectivă comprehensivă asupra regimului eolian de pe teritoriul României, insistând asupra caracteristicilor și tendințelor manifestate în ultimii 50 ani, precum și a influenței Oscilației Nord-Atlantice, utilizând în acest scop diverse categorii de date adunate de la 167 stații meteorologice. Pe baza mediilor multianuale înregistrate și a corelației puternice ($r = 0.87$) stabilite între altitudine și viteza vântului din spațiile interfluviale a fost compusă harta vitezelor medii anuale. Regiunile cu cel mai ridicat potențial eolian sunt Carpații (7-10 m/s pe creste și 3-7 m/s pe versanți și văi) și Litoralul Mării Negre (5-7 m/s); de altfel, cele două regiuni dezvoltă valori extreme ale turbulenței eoliene – maxime în Carpați și minime în unitățile litorale. Interfluviile din Podișul Moldovei și Podișul Dobrogei precum și Bărăganul Nordic prezintă condiții ideale (vânturi medii și puternice și turbulențe scăzute) pentru dezvoltarea parcurilor eoliene. La nivel național, luna cu cea mai mare viteză medie este martie, iar august luna cea mai calmă. În premieră, a fost concepută la scară națională harta direcției rezultante a vântului (DRV), care exprimă atât orientarea rezultantă a vântului cât și intensitatea acesteia (intensitate mare = variabilitate direcțională mică). În ceea ce privește influența NAO se remarcă o corelație negativă între indicii acesteia și viteza vântului (viteza medie, frecvența și intensitatea furtunilor) la nivel multianual și multidecadal. Coeficienții de corelație prezintă valori ridicate în regiunile extracarpatică și valori mici sau medii în cele intracarpatică. Fazele pozitive/negative ale NAO, asociate cu intensitatea scăzută/ridicată a ciclogenezelor din zona mediteraneană, induc viteze mari/reduce ale vântului în acord cu frecvența scăzută/ridicată a ciclonilor care traversează România.

Cuvinte-cheie: *regimul vântului, influența nord-atlantică, direcția vântului*

Introduction

North Atlantic Oscillation (NAO) is one of the most important modes of large scale variability in the Northern Hemisphere. It occurs especially during boreal winter (when the atmosphere is dynamically the most active), while the NAO summer signal is quite null. The phenomenon has been defined as an anomaly in the sea level air pressure (SLP) distribution with centers of action near the Icelandic low and the Azores (Van Loon and Rogers, 1978). Conventionally, the index of the NAO is based on the difference of the standardized sea level pressure (SLP) anomaly measured at Lisbon, Portugal and at Stykkisholmur, Iceland (Hurrell, 1995).

Research studies revealed that changes in the mean air flow and storminess associated with swings in the NAO are reflected in large-amplitude patterns in the anomalies of temperatures, precipitation, heat fluxes and wind speed in the Atlantic-European region (Kushnir, 1994; Hurrell, 1995; Thompson and Wallace, 1998, 2000; Bojariu and Giorgi, 2005).

When the NAO index is positive, enhanced westerly flow across the North Atlantic during winter moves relatively warm (and moist) maritime air over much of Europe, enhanced precipitation over northern Europe associated with less precipitation over central and southern Europe occurs (Hurrell, 1995; Hurrell and van Loon, 1997; Rambu et al., 2002; Bojariu and Gimeno, 2003). A reverse situation occurs during the negative phase of the NAO.

In Romania, positive thermal anomalies are associated with positive NAO phases due to the prevalence of zonal circulation over the Northern and Central Europe and negative thermal anomalies over Romania are associated with negative NAO phases. The winter precipitation pattern over Romania is indirectly related to the NAO (Bojariu and Paliu, 2001; Zaharia et al., 2002).

The study of NAO projection on temperature and precipitation regime in Romania show that the winter NAO related signal is stronger in the extra-Carpathian regions, due to the orographic effects imposed by the Carpathian Mountains on the atmospheric flow (Bojariu and Paliu, 2001).

Previous works of the authors highlight the NAO control upon the climate and the morphodynamics of the Romanian Coast (Vespremeanu-Stroe et al., 2007; Vespremeanu-Stroe and Tățui, 2011).

All the papers which refer to the NAO show that this phenomenon influences all climatic parameters, but there are no references regarding a clear mark of the NAO signal upon the wind regime. The goal of this paper is to reveal the main characteristics and trends of the Romanian wind regime (especially wind speed) and to investigate the NAO projection on it.

Data and Methods

Data

The primary data used in this study consist in monthly and annual means (115 meteorological stations) of wind speed, sums of winter storms (69 stations) and annual means of wind direction (167 stations). In addition, we used the Hurrell's NAO index (Hurrell, 1995). All wind and NAO data covered January 1961 – December 2003 interval.

Methodology

Data used for graphics were normalized extracting from each value the long-term mean and dividing it to standard deviation. The wind speed normalized anomalies were smoothed with a three-year running mean filter to obtain a multi-annual component of the series.

The linkage between NAO variability and Romanian wind regime is identified computing the correlation coefficients between NAO index and wind data and by mapping them. The spatial distribution of wind data and of the correlation coefficients was computed using ArcGIS 3.2, cokriging interpolation method and the wind speed – altitude linear regression (Fig. 1). For storm analysis, we selected wind speed values higher than 10 m/s and 15 m/s, for minimum 6 hour interval. Concerning the wind direction analysis, our interest limited to western circulation frequency expressed by West/East (W/E) ratio which was obtained by dividing the sum frequencies of western winds (from NNW to SSW) to eastern winds (NNE-SSE).




Fig. 1 The influence of altitude on the mean wind speed (wind-exposed areas)

Results

Mean wind speed and storm frequency

The distribution of the mean wind speed over the Romanian territory revealed the influence of the Carpathian chain on the regional response to the large scale atmospheric circulation. The South-Eastern Carpathians represent a roughness element

for the atmospheric flow, acting as a complex barrier (Ion-Bordei, 1988).

In order to quantify the effects of altitude upon wind speed, a correlation between altitude and mean wind speed was computed for 10 mountain meteorological stations placed in wind-exposed areas at altitudes higher than 1200 m (Fig. 1). The resulting correlation coefficient (0.85) indicates a strong linkage; the regression formula was further applied for all mean wind speed data used on the Romanian territory (Fig. 2).




Fig. 2 The mean wind speed distribution over the Romanian territory

As a consequence, the regions between the Carpathians branches (Ardeal) and the intra-mountain and sub-mountain depressions present small annual wind speed values (1-2 m/s), moderate values (2-4 m/s) for the extra-Carpathians regions (Moldova, Muntenia, Oltenia, Dobrogea, Banat, Crișana) while the mountain wind-exposed areas have the highest values (5-9.3 m/s). Another region with relative high wind speed values (4.2-6.95 m/s) is the Romanian Black Sea coast due to the complex connections between the Atlantic storm track, the cyclogenesis activity over the Mediterranean area and the atmospheric circulation over Black Sea (Fig. 2). For the eolian energy, the most suitable locations have to record high speeds and low turbulences

which are rare in Carpathians (especially restricted to wide plateaus: Mt. Mic, Godeanu, Semenic) but frequent in the coastal zone and on the exposed areas of the Moldavian and Dobrogea tablelands or in Bărăgan.

In order to quantify the NAO projection on wind speed pattern over the Romanian territory, we computed the correlation coefficients between Hurrell's NAO index and mean wind speed. For the Romanian territory, the positive wind speed anomalies are associated with negative NAO phases (Fig. 3A). The correlation coefficient (-0.52) is characteristic for the entire Romanian area in the context of a general decreasing trend from large territories to small regions (Fig. 3B, Fig. 4A).




Fig. 3 Time evolution of mean wind speed and Hurrell's NAO index (A) and correlation coefficients between NAOI and wind speed (B) over the Romanian territory

The correlation pattern for wind speed anomalies (Fig. 4A) reveals a strong NAO related signal in extra-Carpathian areas due to an enhanced (diminished) frequency of Mediterranean and arctic cyclones and westerly air flows reaching these regions during negative (positive) NAO phases. The barrier role of the Carpathians against these complex influences determines small correlation coefficients between NAO index and wind speed anomalies in intra-mountain areas and even inverse values in some depressions. A strong NAO signal is found on the Romanian Black Sea coast due to the cyclogenesis activity over the Mediterranean Sea with complex influences upon Black Sea basin. Generally, the same pattern of correlation coefficients spatial distribution occurs for storm frequency and NAO index (Fig. 4B), with a better and more homogeneous distribution in the south-eastern part of Romania, which is an open territory for the Siberian Anticyclone activity (strong winter winds).

Storm distribution during the last four decades of the 20th century show an important shifting at the end of the 70's and the beginning of the 80's, which

is synchronous with NAO signal (Fig. 5). A prominent contrast between a very active interval (1961-1978) and a relatively quiet period with low variability (1979-2003) is noticeable. Temporal evolution of storm activity is in good agreement, but negatively correlated with Hurrell's NAO index.




Fig. 4 Time evolution of storm incidence and Hurrell's NAO index

Another issue of our study was to find the spatial distribution of the windiest interval (monthly) over the Romanian territory, for a multidecadal scale (Fig.

6). It is obvious that there exists a compact grouping of the areas with the same windy intervals controlled by the position of the Carpathians and the west-east distribution. In the Carpathians and along the Black Sea coast, the regular winter (December-February) is the windiest period, while in the mountainous and sub-mountainous depressions and in the limitrophe regions of the Carpathians (e.g. the

Subcarpathians), the windiest interval corresponds to the medium and late spring (April-May). The eastern part of Romania (Dobrogea, Moldavian Plateau, Bărăgan) is affected by strong winds at the end of winter (February-March). The central and the western parts record the highest wind speed at the beginning of the spring: March-April.




Fig. 5 The NAO patterns of correlation coefficients for mean wind speed (A) and storm frequency (B) over the Romanian territory




Fig. 6 The windiest interval distribution over the Romanian territory

Regarding the mean monthly wind speed, January – April is the most active period over the entire Romanian territory, with the highest values of wind speed (3 – 3,3 m/s) and the maximum means in March, while August is the most calm month of the year. The greatest variability of wind regime is registered in November – March interval, with maximum in January and February (winter conditions) and minimum in May – September under summer conditions (Fig. 7).




Fig. 7 The mean monthly wind speed in Romania (115 meteo stations). Gray columns represent the mean wind speed and the black overimposed small columns the standard deviation

Resultant wind direction

Based on the directional wind speed and frequency, we computed the direction and magnitude of the vectors representing the Resultant Wind Direction (RWD) for 167 meteorological stations spread over the Romanian territory (Fig. 8). The highest magnitudes of RWD (expressed in vector-units: vu) exceed the value of 200 vu and are representative for the meteorological stations placed on the alpine plateaus of the Romanian Carpathians. The general directional trend for these vectors is eastward, sometimes with a south-eastern component (especially for the Southern Carpathians). As expected, the meteo stations situated in the intra-mountainous areas (valleys, corridors, depressions) show a directional distribution of the RWD vectors in accordance with their general orientation.

The directional distribution of the vectors shows some discrepancies between W/E and N/S ratios. Thus, 75% of the total vectors are oriented to the East (from 0 to 180 degrees) and only 25% to the West (from 180 to 360 degrees), while 69% are oriented to the South (from 90 to 270 degrees) and only 31% to the North (from 270 to 90 degrees).

Moreover, the strongest resultant directions (> 50 vu) are recorded in the most cases (95%) on eastward and southward directions. Also, the average direction of the vectors is 141 degrees (SE). These results give a very good picture of the general airflows which pass over the Romanian territory and which are oriented to the east and south. A marked pattern is the directional grouping of the vectors for the meteo stations placed in the same region of the country. Thus, we separated the following situations: i) the vectors for the Eastern part of Romania which are oriented mainly to the SE (for Moldova) and S (for Dobrogea), according to the orographic effect of the Eastern Carpathians and the influence of the Black Sea; ii) the southern part of the

country (Muntenia and Oltenia) is dominated by southward air flows over the Subcarpathians and Getic Plateau, due to the influence of the Southern Carpathians. The most part of the low areas (plain level) are affected by northeasterly winds, while the imprint of the Danube corridor gives a south-eastward component, especially for the meteo stations situated on the Danube River; iii) the vectors from the central part of Romania (Transilvania) are south-eastern oriented as a result of the air flows from West which dominate this region; iv) the vectors from the western part of the country do not present a general direction trend, possible due to the complex air flows which pass this area from North, South and West.




Fig. 8 The resultant wind direction over the Romanian territory

The NAO and wind direction

As NAO controls the mass circulation pattern over the Europe, we assessed the NAO role in western circulation strengthening or weakening over the Romanian territory. The temporal pattern of W/E wind frequency ratio during the last four decades indicates an intensification of western circulation in the last two decades (Fig. 9A). In this

case a direct proportionality relation is established between the western circulation frequency and NAO. Positive/negative anomalies of western circulation frequency over the Romanian territory are associated with positive/negative NAO phases (positive statistical dependency) with a good correlation coefficient (0.69) based on wind data from 122 meteorological stations (Fig. 9B).




Fig. 9 Time evolution of W/E wind frequency ratio and NAOI (A) and correlation coefficients between NAOI and western circulation (B) over the Romanian territory

The correlation pattern for W/E wind frequency indicates a strong positive dependent NAO signal in extra-Carpathian areas (especially in south and south-east). The arch configuration of the Carpathians standing against large scale circulation determines small correlation coefficients in the center and eastern part of Romania and very small correlation values in intra-mountain and sub-mountain areas.

Conclusion

It is widely accepted that NAO controls the overall distribution of temperatures and precipitations in Europe (van Loon and Rogers, 1978; Hurrell and van Loon, 1997; Bojariu and Paliu, 2001; Râmbu et al., 2002; Zaharia et al., 2002); our results indicate that the wind regime over the Romanian territory is also strongly related to NAO phases (Fig.4, Fig.5).

Over the Romanian territory, the position of the Carpathians mountains induces a high variability for the spatial distribution of the eolian parameters characteristics and influences the local response to the NAO variability. Thus, the extra (intra-) Carpathian regions are characterized by high (medium to low) correlation coefficients between the NAO index and wind anomalies.

The correlation coefficients point out that positive wind speed and storm occurrence anomalies are associated with negative NAO phases (negative statistical correlation), while positive W/E wind frequency variability is correlated with positive NAO phases (positive statistical dependency).

The positive (negative) NAO associated with low (high) cyclogenesis over the Mediterranean area

induce low (high) winds over the Romanian territory due to diminished (enhanced) frequency of cyclones reaching Romanian regions.

Acknowledgements

Special thanks are addressed to Florin Iosub for helping us mapping the Resultant Wind Direction values.

References

- Bojariu, R., Paliu, D.M., 2001. North Atlantic Oscillation projection on Romanian climate fluctuations in the cold season, in *Brunet and Lopez* (Eds.), *Detecting and Modelling Regional Climate Change and Associated Impacts*.
- Bojariu, R., Gimeno, L., 2003. Predictability and numerical modeling of the North Atlantic Oscillation, *Earth – Science reviews*, 63(1-2), 145-168.
- Bojariu, R., Giorgi, F., 2005. The North Atlantic Oscillation signal in a regional climate simulation for the European region, *Tellus*, 57A,1-13.
- Hurrell, J.W. 1995. Decadal trends in the North Atlantic Oscillation: regional temperature and precipitation. *Science*. 269, 676-679.
- Hurrell, J.W., van Loon, H., 1997. Decadal variations in climate associated with the North Atlantic oscillation. *Climatic change*, 36, 301-306.
- Ion-Bordei, N., 1988. *Meteoclimatic phenomena induced by the Carpathian configuration in the Romanian Plain*. Ed. Academiei, 174 pp (in Romanian).
- Kushnir, Y., 1994. Interdecadal variations in the North Atlantic sea surface temperature and associated atmospheric conditions. *Journal of Climate*, 7, 141-157.