Forum Geografic - Studii și cercetări de geografie și protecția mediului (FG - S.C.G.P.M.)

The Journal Forum geografic. Studii de geografie și protecția mediului (Geographical Phorum – Geographical Studies and Environment Protection Research) was founded in 2002, and it seeks to publish high quality research in the domains of geography, environment protection and other related domains. It addresses a range of issues, such as geomorphology, pedology, climatology, hydrology, human geography and environment. Its content is directed to a broad audience, including both academics and policymakers. The papers selected for publication in the journal are subject to a review process using references from universities worldwide.

The journal is currently indexed by the following databases: SCOPUS, DOAJ, EBSCO, ERIH PLUS, Index Copernicus, Scipio, CrossRef, Ulrichsweb, IGU, Google Scholar, WorldCat, ProQuest, ZDB, CNCSIS, DRJI, citefactor.org journals indexing, InfoBase index, Eurasian Scientific Journal Index, ResearchBib.

Editor-in-chef: Sandu BOENGIU, Physical Geography, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Liliana POPESCU, Human Geography, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Executive editor: Oana Mititelu Ionuș, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Tehnical editor: Cristiana Vîlcea, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Editorial Advisory Board:

Iuliana ARMAŞ, University of Bucharest, Romania	Oana MITITELU IONUȘ, University of Craiova, Romania
Lucian BADEA, The Institute of Geography, The Romanian Academy	Emil MARINESCU, University of Craiova, Romania
Dan BĂLTEANU, The Institute of Geography, The Romanian Academy	Mirela MAZILU, University of Craiova, Romania
Zeljko BJELJAC Geographical Institute Jovan Cvijić, Serbia	Ciprian MĂRGĂRINT, Al.I.Cuza University, Iași, Romania
Lucian BLAGA, University of Oradea, Romania	Alexandru NEDELEA, University of Bucharest, Romania
Sandu BOENGIU, University of Craiova, Romania	Amalia NIȚĂ, University of Craiova, Romania
Léon BRENIG, University of Brussels, Belgium	Zvi Yehoshua OFFER, Ben-Gurion University, Israel
Manola BRUNET, University of Tarragona, Spain	Aurel PERȘOIU, Emil Racovita Institute of Speleology, Romanian Academy
Sorin CHEVAL, Henri Coandă Air Force Academy, Brașov, Romania	Dănuț PETREA, University of Babeș-Bolyai, Cluj-Napoca,Romania
Laura COMĂNESCU, University of Bucharest, Romania	Liliana POPESCU, University of Craiova, Romania
Lóczy DÉNES, University of Pécs, Hungary	Kumar Rai PRAVEEN, Banaras Hindu University, India
Philippe DONDON, Ecole Nationale Superieure E.I.T.M.M. de Bordeaux/ Institut	Milan RADOVANOVIĆ, Geographical Institute Jovan Cvijić, Serbia
Polytechnique de Bordeaux, France	Khaiwal RAVINDRA, School of Public Health, PGIMER, Chandigarh, India
Monica DUMUTRAȘCU, The Institute of Geography, The Romanian Academy	Igor SIRODOEV, Universitatea "Ovidius" din Constanța, Romania
Dan DUMITRIU, Al.I.Cuza University, Iași, Romania	Daniel SIMULESCU, National Agency for Protected Natural Areas, Romania
Recep EFE, Balikesir University, Turkey	Fábián Ákos SZABOLCS, University of Pécs, Hungary
Robert FOVELL, University of California, USA	Gheorghe ŞERBAN, University of Babeş-Bolyai, Cluj-Napoca,Romania
Teodoro GEORGIADIS, Istitute of Biometeorology (IBIMET), Bologna, Italy	Cristina ȘOȘEA, University of Craiova, Romania
Adrian GROZAVU, Al.I.Cuza University, Iași, Romania	Titus MAN, University of Babeș-Bolyai, Cluj-Napoca,Romania
Nelly HRISTOVA, St. Kliment Ohridsky University of Sofia, Bulgaria	Magdy TORAB, Alexandria University, Egypt
Ioan IANOŞ, University of Bucharest, Romania	Marcel TÖRÖK – OANCE, West University of Timişoara, Romania
Corneliu IAȚU, Al.I.Cuza University, Iași, Romania	Cristiana VÎLCEA, University of Craiova, Romania
Alexandru ILIEŞ, University of Oradea, Romania	Alina VLĂDUȚ, University of Craiova, Romania
Mihaela LICURICI, University of Craiova, Romania	Nenad ŽIVKOVIĆ, Belgrade University, Serbia
Dumitru MIHĂILĂ, Ștefan cel Mare University, Romania	Martina ZELENAKOVA, Technical University of Kosice, Slovakia
Gabriel MINEA, National Institute of Hydrology and Water Management, Romania	Zbigniew ZWOLIŃSKI, Adam Mickiewicz University (Poznan), Poland

Associate Editors:

Slavoljub DRAGIĆEVIĆ, Faculty of Geography, Belgrade University, Studentski trg 3/3, Belgrade, Serbia Vesna LUKIĆ, Demographic Research Centre, Institute of Social Sciences, Kraljice Natalije 45, Belgrade, Serbia Nina NIKOLOVA, Faculty of Geology and Geography, "*St. Kliment Ohridsky*" University of Sofia, Tzar Osvoboditel Blvd. 15, Sofia, Bulgaria Gabriel MINEA, National Institute of Hydrology and Water Management, *Romania*

Assistant Editors: Daniel SIMULESCU, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Founding Editors: Boengiu S., Marinescu E., Pleniceanu V., Tomescu V., Enache C.

Cover photo: Metaliferi Mountains, Alba county (by Alina Vlăduț)

For instructions for authors, subscription and all other information please visit our website <u>http://forumgeografic.ro</u>

before submitting any papers please select the section Publishing rules from the About page and read thoroughly the submission instructions for authors

ISSN 1583-1523 (print) ISSN 2067-4635 (online) DOI prefix: 10.5775

Article submission

In order to disseminate the research results in the field, researchers, scholars and professionals are welcome to submit an electronic version of the manuscript (in Microsoft Office Word format) to the editorial office (forum.geografic@gmail.com).

Submission requirements: The submission of an article for publication in our journal implies that the article has not been published before, nor it is being considered for publication in other journals. Authors are responsible for the content and the originality of their contributions. In order to be published, articles must be thoroughly researched and referenced.

IMPORTANT: All papers must be submitted in electronic format, only in English language.

Copyright statement

By submitting a scientific work to *Forum geografic* the submitters agree to declare the following:

- the submitted work belongs exclusively to the declared authors;
- the submitted work represents original scientific research;
- the submitted work has not been published or submitted for publishing to another journal;
- if the submitted work is published or selected for publishing in *Forum geografic*, the authors waive any patrimonial claims derived from their authorship for the submitted work; the authors retain the moral rights for their submitted work, as granted under the Romanian applicable law; also, the authors agree to refrain from ulterior submitting of the work to other journals.

The submitters agree to be solely held accountable in case of breaching the above terms and to defend the representatives of **Forum geografic** in the event of a lawsuit related to the submitted work.

When submitting a paper, the authors are required to print, fill and send a scanned copy of this declaration.

Privacy statement

The submitted personal data, such as names or email addresses, are used only for the declared purpose of the *Forum geografic* journal (publishing original scientific research) and are not available to third parties.

Manuscripts are received at all times. However, in order to have your article published in the current year, the manuscripts must be submitted until the 15^{th} of February for the first issue of the current year and until the 1^{st} of September for the second issue.

Article format

All manuscripts must be edited entirely in English. Articles must include:

- Title
- Author's name(s). For each author you must mention the author's scientific title, his affiliation (institution) and e- mail address;

- Abstract (maximum 300 words);
- Keywords (not more than 5-6 words);
- Acknowledgments (if any);
- Main body of text (structured according to Introduction, Data & Methods, Results & Discussions, Conclusions);
- Illustrations (graphs, diagrams, maps, photos should have indications of their positions in the text and title written in English) must be also submitted in electronic format, preferably in JPG, PNG or BMP format and must be referred to as Figures, which should be numbered with Arabic numbers.
- Tables must be numbered with Arabic numbers and should not repeat data available elsewhere in the text.
- References must be indicated in the text, between brackets and they must include the author's name and the date of the publication (Popescu, 2000). When three or more authors are referred, they will appear in the text as follows: (Popescu et al., 1997). References must be listed in alphabetical order at the end of the text.

The following style sheet is recommended:

• for journals:

Miletić, R., Lukić, V., & Miljanović, D. (2011). Deindustrialization and structural changes in commuting flows in Serbia. *Forum geografic*, X(2), 244-254. doi:10.5775/fg.2067-4635.2011.009.d

for books:

Bran, F.,Marin, D., & Simion, T. (1997). Turismul rural. Modelul european, Editura Economică, București

• for papers from conference proceedings:

Deci, E. L., Ryan, R. M., (1991), A motivational approach to self: Integration in personality. In R. Dienstbier (Ed.), *Nebraska Symposium on Motivation: Vol. 38. Perspectives on motivations* (pp. 237-288). Lincoln: University of Nebraska Press.

Review process

All the manuscripts received by the editors undergo an anonymous peer review process, necessary for assessing the quality of scientific information, the relevance to the field, the appropriateness of scientific writing style, the compliance with the style and technical requirements of our journal, etc. The referees are selected from the national and international members of the editorial and scientific board, as well as from other scholarly or professional experts in the field. The referees assess the article drafts, commenting and making recommendations. This process leads either to acceptation, recommendation for revision, or rejection of the assessed article. Editors reserve the right to make minor editorial changes to the submitted articles, including changes to grammar, punctuation and spelling, as well as article format, but no major alterations will be carried out without the author's approval. Before being published, the author is sent the proof of the manuscript adjusted by editors. If major revisions are necessary, articles are returned to the author so that he should make the proper changes. Authors are notified by email about the status of the submitted.



Forum geografic. Studii și cercetări de geografie și protecția mediului
Volume XXII, Issue 1 (June 2023), pp. 5-15; DOI: 10.5775/fg.2023.090.i
© 2023 The Author(s). Published by Forum geografic.
Open Access article.
Open Access article.
Creative Commons Attribution-NonCommercial-NoDerivatives License.

A statistical analysis of temperature and precipitation in Belgrade, Serbia (1961-2020)

Nina ČEGAR^{1,*}, Uroš DURLEVIĆ¹, Milica DOBRIĆ¹, Sandra VUKAŠINOVIĆ¹

¹ Faculty of Geography, University of Belgrade, Studentski Trg 3/III, Belgrade, Serbia * Corresponding author. nina.cegar@gef.bg.ac.rs

Received on 18-10-2022, reviewed on 27-12-2022, accepted on 27-03-2023

Abstract

The most significant effects of climate change are related to temperature and precipitation. Changes in precipitation and temperature patterns affect water resource capacity, agricultural activities, global biodiversity and emergency management. In this study, using two statistical models (Mann-Kendall and Pettit's test), climate data for the territory of Belgrade in the time interval 1961-2020 were processed. Trends were analyzed: average annual temperatures, average monthly temperatures, annual and monthly precipitation amounts. The aim of the research is to determine the statistical significance of the increasing/decreasing temperature and precipitation trends on a monthly and annual basis. Based on the obtained results, it was determined that the changing point of the annual temperature is in 1997, while a statistically significant increasing trend was observed in all months except November. The increasing trend recorded by the precipitation amount is 0.345 mm/year, but without statistical significance. The most significant results corresponding to July and October. **Keywords:** *Trend, Mann-Kendall test, Pettit's test, precipitation, air temperature, City of Belgrade*

Introduction

Climate change is reflected in large fluctuations in climate averages that persist for decades or even longer periods. Although climate change occurs on a global scale, impacts often vary from region to region. In the 20th century numerous climatic anomalies were observed- poor monsoon years around the turn of the century, rapid Arctic warming into the 1920s, the Dust Bowl drought and heat waves in North America in the 1930s and drought in Australia, and cold winters and hot summers in Europe in the 1940s. These anomalous events occurred during a period of strong global-scale warming, which can be attributed to a combination of external forcing (particularly, greenhouse gas increases, combined with a hiatus in volcanic events) and internal decadal variability. During the 20th century, in most European regions, a temperature increase was observed (Trajkovic & Kolakovic, 2009; Gocic & Trajkovic, 2013; Hegerl et al., 2018). Climate reconstructions show that summer air temperatures in Europe in period 1986-2015 have been the warmest for at least 2000 years, and that they lie significantly outside the range of natural variability (Luterbacher et al., 2016). Based on data from 210 weather stations on the territory of Europe, temperature increase amounts 0.051°C/year in the period 1985-2020. The highest increase was recorded in spring, the lowest in autumn (Twardosz et Vyshkvarkova and Sukhonos (2022) al., 2021). investigating the spatial distribution of temperature and precipitation extremes in Eastern Europe, conclude that there are positive and statistically significant trends for

warm extremes in all seasons, with maximum values in the winter season. In contrast, negative trends were obtained for cold extremes.

Each of the last four decades has been successively warmer than any decade that preceded it since 1850. Global surface temperature in the first two decades of the 21st century (2001–2020) was 0.99 (0.84 to 1.10) °C higher than 1850–1900. Global surface temperature was 1.09 [0.95 to 1.20] °C higher in 2011-2020 than 1850–1900, with larger increases over land (1.59 [1.34 to 1.83] °C) than over the ocean (0.88 [0.68 to 1.01] °C). Hot extremes (including heatwaves) have become more frequent and more intense across most land regions since the 1950s, while cold extremes (including cold waves) have become less frequent and less severe. Globally averaged precipitation over land has likely increased since 1950, with a faster rate of increase since the 1980s. Mid-latitude storm tracks have likely shifted poleward in both hemispheres since the 1980s, with marked seasonality in trends (IPCC, 2021) According to NOAA (2022), all years in the period 2013-2021 belongs to the category of ten warmest years since measurements were made. Since 1981, the average increase in global temperature is 0.18°C per decade. In 2021 the western parts of the USA, southern South America, northern Africa and the Middle East were characterized significantly by below-average precipitation. A significant increase in annual precipitation during 2021 was observed in northern South America, eastern Europe, western and eastern parts of the Australian coast, and South Asia. Across Southern Europe, below-average rainfall was recorded in July. Dry conditions with high temperatures contributed to the spread of fires in southern and southwestern Turkey, as well as in north-eastern Spain. In Spain, during July 2021, only 57% of the average amount of July precipitation fell. However, parts of western Europe faced torrential rainfall that caused flooding between July 13 and 15. Some locations received at least twice their normal monthly precipitation in only three days. According to preliminary reports, over 140 deaths are attributed to the floods.

Branković et al. (2013) determined for Croatia that the positive trends of the average annual temperature are caused by a statistically significant increasing trend in the summer months. The highest trend values were obtained for summer, followed by winter and spring. As the research conducted by Gocic & Trajkovic (2013) shows, there is a significant increase in temperatures in the northern, central and eastern parts of Serbia. A constant annual increase in maximum daily temperatures from 1974-2003 in Belgrade and Niš was observed by Milanovic et al. (2014). The results of the Pettit's test of average annual temperatures in the area of Sokobanja in the period 1946–2012 indicate that 1991 was a changing point in the temperature increase (Radivojević et al. 2015).

Luković et al. (2014) investigated the trend of precipitation in Serbia in the period 1961-2009. No significant trend of total annual precipitation has been detected for the whole territory of Serbia. Very slight tendencies toward drier conditions on a seasonal scale during winter and spring and wetter conditions during autumn. An increase in autumn amounts of precipitation over central parts of Serbia is a result of a significant increase in October precipitation. The results obtained by Malinovic-Milicevic et al. (2016) indicate a significant increasing trend in the amount and intensity of precipitation in Serbia during autumn. Bandyopadhyay & Perveen (2006) and Gajbhiye et al. (2015) point out that changes in the amount and frequency of precipitation, as a result of climate change, directly affect water flows, runoff, groundwater reserves and soil moisture. Determining the change in rainfall patterns is crucial for any kind of development program and planning that includes effective management of water resources, but also the food production sector.

The objectives of this study are to investigate the temperature and precipitation trends at annual and monthly level in Belgrade in the period 1961-2020 using the Mann-Kendall test. The Pettit's test was used to determine if there were changing point years of both parameters. The paper will also present a comparative analysis of the values of selected parameters in two climatological periods 1961-1990 and 1991-2020 and inter–annual variability of precipitation.

Study area

The study area is Belgrade, the capital of the Republic of Serbia (Figure 1). It covers an area of 360

km², while the wider city area covers 3 222 km². It is located in the Pannonian Plain, on the southern edge of the Pannonian Basin. It lies at the confluence of two large European rivers - the Sava and the Danube rivers, and partly belongs to the Balkan Peninsula, partly to Central Europe (Drazic et al., 2014).



Figure 1: Geographic position of Belgrade in the Republic of Serbia

The study area map was obtained in geographic information systems, which play a major role in the analysis of phenomena and processes in the environment (Ćurić et al., 2022)

Belgrade is located in the zone of moderate continental climate (Meteologos, 2012). The climate of Belgrade is characterized by a great variability of the meteorological elements, which is caused by the circulation of air masses with different physical features. For the territory of the city, intrusions of polar and tropical air are characteristic. Relief has a significant influence on the microclimate of the capital of Serbia. It is characterized by the vast Pannonian plain in the north and the undulating surfaces of northern Šumadija and Avala Mountain (511 m) in the south. The Sava and the Danube rivers, at the confluence of which the city is located, are also significant factors.

Warm and cold periods often occur in all seasons. Cold and moist air are intruding from the northwest and west, and heavy precipitation is brought by the cyclones from the western Mediterranean Sea, which move eastward through the Sava and Danube valleys. Northern intrusions across the Pannonian plain cause a noticeable drop in temperature, while intrusions of cold air from the northeast, from the Carpathians, in winter, cause cold, windy and mostly dry weather. At the beginning of May, the intrusions of moist and cold air appear. Local showers and thunderstorms are frequent in May and June. Shorter and longer dry periods are characteristic in summer (Unkašević, 1994). Dominant winds are košava, west and north-west winds (Meteoblue, n.d.).

Data and methods

The data come from the Republic Hydrometeological Service of Serbia (RHMSS) measured at the weather station Belgrade (Serbia). The coordinates of the meteorological station are 44°48 'N and 20°28 'E and it is located at an altitude of 132m (Hidmet, n.d.). The paper analyzed mean monthly and mean annual air temperatures, monthly and annual precipitation during two climatological periods - 1961-1990 and 1991-2020.

The Mann-Kendall test (Mann 1945, Kendall, 1975) was used for trend analysis. The test belongs to the group of non-parametric statistical trends that treats series that are not normally distributed, and its use is based on multi-year data series. Mann–Kendall is commonly used to detect monotonic trends in climatological and hydrological research. The Z statistic was used to test significance. A positive Z value indicates an increasing trend, while a negative Z value indicates a negative trend (Helsel & Hirsch, 2002; Westra et al., 2013; Pohlert, 2015). The null hypothesis of the Mann-Kendall test is based on the assumption that there is no monotonic trend in the time series. Control statistic is used to test the null hypothesis (Kendall, 1975).

The Mann-Kendall (S) Statistic is computed as follows (Kendall, 1962):

 $S = \sum_{i=1}^{n-1} k \sum_{j=i+1}^{n} sign(Tj - Ti),$

where Ti and Tj are the values of sequence i, j; n is the length of the time series and:

Sign
$$(Tj - Ti) = \begin{cases} 1 \ if \ Tj - Ti > 0 \\ 0 \ if \ Tj - Ti = 0 \\ -1 \ if \ Tj - Ti < 0. \end{cases}$$

If the Z value (significance level) is greater than 1.96 (which corresponds to the significance threshold of 0.05), we conclude that there is a monotonic trend in the time series, and otherwise, if the value is lower, the trend does not exist (Stojković et al., 2014). Sen's estimate of the slope (Sens's estimate) shows us the estimate of the slope of the linear trend and the average value of changes in a unit of time (Salmi et. al., 2002; Ahmad et. al., 2015).

In order to detect significant changes in time series of climatological data, the Pettit's test was used. The Petit test also belongs to the group of non-parametric tests and is most often used to detect sudden changes (points of change) in climatological data. The Pettit test is a method that discover a significant change in the mean value of a time series when the exact time of the change is unknown. According to Pettitt's test, if $x_1, x_2, x_3, ... x_n$ is a series of observed data which has a change point at t in such a way that $x_1, x_2, ..., x_t$ has a distribution function F1(x) which is different from the distribution function F2(x) of the second part of the series $x_{t+1}, x_{t+2}, x_{t+3}, ..., x_n$. The non-parametric test statistics Ut for this test may be described as follows (Pettit, 1979):

$$Ut = \sum_{i=2}^{n} r \sum_{j=1}^{i-1} sign(Xi - Xj),$$

$$sign (Xi - Xj) = \begin{cases} 1 \ if (Xi - Xj) \\ 0 \ if \ (Xi - Xj) \\ -1 \ if \ (Xi - Xj). \end{cases}$$

The test statistic counts the number of times that a member of the first sample exceeds a member of the second sample. The null hypothesis of Pettitt's test is the absence of a changing point. The test statistic (K) may be described as:

K = Max |Ut|

When the value of the test statistic is less than the chosen confidence interval, the null hypothesis is rejected and there is no distinct change point in the time series (Jaiswal et al., 2015).

The inter–annual variability of precipitation was obtained by subtracting the annual total of each year from the mean annual total over the study period and dividing the difference by the standard deviation of the series. Thus, obtaining a new series in which the mean equals to zero and the variance to 1. The obtained values enable the division of all years into three categories as follows (Kutiel et. al., 2015):

Dry (D) when z < -0.5Normal (N) when $-0.5 \le z \le 0.5$ Wet (W) when $0.5 \le z$

Results and Discussion

Temperature

The average annual air temperature in Belgrade in the period 1961–2020 is 12.5 °C. The warmest month is July with an average temperature of 22.8 °C, and the coldest is January with an average temperature of 1.1 °C (Table 1). The warmest year in the observed period was 2019 (14.8 °C), and the coldest year was 1980 (10.8 °C).

The mean annual air temperature in Belgrade shows an increasing trend of 0.042 °C/year with a significance level of 0.001. There is an increasing trend of mean monthly temperatures. Only in November the temperature increase is not statistically significant. There is a distinct trend of growth in the summer months (June, July, August) – 0.046 °C/year, 0.06 °C /year and 0.075 °C /year respectively (Table 2, Figure 2).

Month	Ι	II	III	IV	V	VI	VII	VIII	IX	Х	XI	XII	Annual
T _A (°C)	1.1	3.3	7.7	13.0	17.7	21.0	22.8	22.6	18.1	12.9	7.5	2.7	12.5
Same Bruchlis Hudensets and sized Camies of Carbin Climets Verslands 1001 2020													

Source: Republic Hydrometeorological Service of Serbia, Climate Yearbooks 1961-2020

Table 2: Mann-Kendall test results for average annual and average monthly air temperatures inBelgrade (1961-2020)

Month	Trend (Z)	Sen's estimate (B)	Level of significance (a)
January	2.45	0.045	*
February	1.88	0.048	+
March	2.44	0.046	*
April	2.98	0.043	**
May	1.95	0.026	+
June	3.68	0.046	***
July	5.20	0.060	***
August	4.95	0.075	***
September	1.89	0.027	+
October	2.09	0.032	*
November	1.29	0.023	
December	2.67	0.041	**
Year	6.07	0.042	***

*** - significance level 0,001; **- significance level 0,01; *- significance level 0,05; +- significance level 0,1



Figure 2: Trend of average annual air temperature and average temperatures of summer months in Belgrade (1961 - 2020)



Figure 3: Pettit's test results for average annual temperatures in Belgrade (1961-2020)

The statistically significant changing point year according to the Pettit's test is 1997 (p=0.0001). The period after 1997 recorded an average annual air temperature of 13.4 °C compared to 11.9 °C in the period from 1961 to 1997 (Figure 3).





Since the statistically most significant increasing trends were recorded in the summer months, it was investigated whether there is a changing point for each of them. The average June temperature in the sixtyyear period is 21.7 °C, and in 1990 there was a change (increase) in the average monthly temperature from 20.1 °C to 21.9 °C (p=0.0001). The average monthly temperature in July in Belgrade in the observed period is 21.5 °C. The statistically significant year in the average temperature change is 1986 (p p=0.0001). Until the mentioned year, the average temperature was 21.5 °C, so it corresponds to the sixty-year average. After 1986, the average July temperature in Belgrade rose to 23.8 °C. Since 1991 (p=0.0001), the average air temperature in August has increased by 2.5 °C; from 21.3 °C to 23.8 °C (Figure 4).

Table 3: Results of Pettit's test for average annual and monthly temperatures in Belgrade (1961-2020)

Parameters	Level of significance (p)	Changing point (year)	Change (°C)
Average annual	0.0001	1997	+1.483
temperature			
Average			
temperature in	0.0001	1990	+1.845
June			
Average			
temperature in	0.0001	1986	+2.292
July			
Average			
temperature in	0.0001	1991	+2.505
August			

The results indicate that the temperature in Belgrade is inevitably rising. An increasing trend without statistical significance was only observed in November, while in all other months the results showed a statical significant increasing trend. Pešić and Milovanović (2016) used the same methodology to determine temperature changes in the area of Sumadija (Serbia) in the period 1961-2010. They obtained a statistically significant increasing trend of annual temperature at almost all stations. The second highest mean annual temperature increase is in Belgrade (0,03°C/year), which is similar to obtained results in this paper (0,04°C/year). In Serbia, in the region of Vojvodina and Kosovo and Metohija positive air temperature trends were also obtained (Gavrilov et al., 2015; Gavrilov et al., 2016; Gavrilov et al., 2018). The average temperatures increasing trend indicates that the warming of the climate system is present in the entire territory of Bosnia and Herzegovina according to the results obtained by Trbić et al. (2017). The warming trend is more pronounced since 1990, the highest values in the period 1961–2015 were obtained in the warmest part of the year and in the area of Banja Luka and Bijeljina. In the region, temperature increase is also dominant in Montenegro (Burić et al., 2014).

Climatic periods	Ι	II	III	IV	v	VI	VII	VII	IX	x	XI	XII	Annual
1961- 1990	0.4	2.8	7.2	12.4	17.2	20.1	21.8	21.4	17.7	12.5	7.0	2.3	11.9
1991- 2020	1.9	3.8	8.3	13.6	18.2	21.9	23.8	23.8	18.5	13.3	8.1	3.0	13.2

Table 4: Average temperature (°C) in B	Igrade during two climatic periods 1961-1990 and
1991-2020	

Source: Republic Hydrometeorological Institute of Serbia, Meteorological yearbooks 1961–2020

Since the data of two climatological periods were used, the data on the average values of air temperature by period - from 1961 to 1990 and from 1991 to 2020 - are presented (Table 4). The average temperatures in Belgrade in the last period are higher during all months compared to the average temperatures in the period from 1961 to 1990. The biggest difference is in the summer months – August (2.4 °C), July (2 °C) and June (1.8 °C). In the same months, trend analysis showed the most significant increases in temperature. In the period from 1961-2010, according to research by Bajat et al. (2014), a significant increase in temperature in the summer period at 22 meteorological stations in Serbia was established.

In July 2012, the highest average monthly temperature (27 °C) in the observed period was recorded. During 2012, which is characterized as extremely hot and dry year, temperatures in Serbia were above 35 °C for more than 50 days.

Temperatures were extremely high in 2007, and precipitation were unevenly distributed (Živanović et al., 2020). The absolute maximum temperature of 43.6 °C in Belgrade was measured on July 24, 2007. On the same date, the highest temperature of 44.9 °C in the Republic of Serbia was measured in Smederevska Palanka (Anđelković, 2007). The largest number of fires (28,548) in the period from 2000 to 2010 occurred in 2007 (Živanović et al., 2020).

Precipitation

The average annual amount of precipitation in Belgrade in the period from 1961 to 2020 is 691.8 mm. On average, at monthly level, the highest amount of precipitation is in June 93 mm and May 71.5 mm, then in September 55 mm. The "driest" month is February with an average of 44 mm of precipitation (Table 5).

Month	Ι	II	III	IV	V	VI	VII	VIII	IX	Χ	XI	XII	Annual
Average precipitation (mm)	48.9	44.0	49.1	55.1	71.5	93.0	66.4	53.2	55.0	47.6	51.9	56.1	691.8
Source: Republic Hydrometeorological Service of Serbia, Meteorological yearbooks, 1961–2020													

Source: Republic Hydrometeorological Service of Serbia, Meteorological yearbooks 1961–2020

Table 6. Mann-Kendall test results f	or annual and monthly j	precipitation in Belgrade (19	61-
2020)			

Month	Trend (Z)	Sen's estimate (B)	Level of significance (a)
January	0.44	0.128	
February	0.59	0.155	
March	0.27	0.074	
April	-1.47	-0.243	
May	1.01	0.277	
June	0.85	0.386	
July	-1.65	-0.416	+
August	0.78	0.208	
September	-0.15	-0.026	
October	2.67	0.592	**
November	-1.39	-0.283	
December	-0.99	-0.283	
Year	0.24	0.345	

*** - level of significance 0,001; **- level of significance 0,01; *- level of significance 0,05; + - level of significance 0,1



Figure 5: Precipitation trend in July and October with Sen's estimate in Belgrade 1961-2020

At the annual level, non-significant increasing trend was observed. In July, there is a decrease in the amount of precipitation (0.416 mm/year) with a significance level of 0.1. Decreasing precipitation trend is also present in April, September, November and December, but they are not statistically significant. The precipitation increase is observed in January, February, March, May, June, August but without statistically significance, while in October the increasing trend (0.592 mm/year) has a level of significance of 0.01 (Table 6, Figure 5). Changing point wasn't detected in the amount of precipitation.

No significant changes in the annual amount of precipitation were obtained in the work. Luković (2013) obtained the same results at the state level in the period 1961-2009, and Gocić and Trajković (2013) didn't found significant rainfall trend at most station in the period 1980-2010. In current paper, the most significant increasing trend was obtained in October (autumn), similar to research by and Gocić and Trajković (2013) and Tošić et. al. (2014) which obtained a significant precipitation increase during autumn and winter.

It is difficult to find similar rainfall results in the literature over a wider area. The area of south-eastern and central Europe recorded either decreases in precipitation from -20 to 0 mm/decade or an increase in precipitation from 0 to 20 mm/decade, whereby the

boundary of change from the negative to the positive trend passes through Serbia to around 22° eastern longitude, but it is a generalized survey (EEA, 2017; Milentijević et al., 2020;). For most parts of Serbia, the mean annual precipitation trend for the period 1961 to 2010 falls in between -5 and +5 and +5 and +15 mm/decade (Milovanović et al., 2017). Popov and Svetozarevich (2021), analyzed annual rainfall trend on 15 weather station in Serbia The least increase in the period 1991-2019 has been observed at the stations of Belgrade, Novi Sad and Zrenjanin. The highest values are reported at Kopaonik station (84.5 mm/decade) and Crni Vrh station (70.1 mm/10 decade).

During the period 1961-2010, a slight positive trend in the annual precipitation was obtained in the eastern part of Croatia, while changes in the annual rainfall wasn't record in Hungary. (Klapwijk et al. 2013, Gajić - Čapka et al.,2015). During the period 1961-2013 some stations in western Romania recorded a statistically significant increase in the mean annual precipitation, on the other hand, the stations in the southwest of Romania recorded a statistically significant decrease in the mean annual precipitation (Croitoru et al. 2016).

The difference in monthly precipitation between the two climate periods (1961-1990 and 1991-2020) varies. A decrease in monthly precipitation is observed in the

winter months (December, January and February) and in March, April and November. The largest decrease in monthly precipitation between the two periods is in April – 7,3 mm (Table 7). The largest increase (14,5 mm) in monthly precipitation in the second thirty-year period is in October.

According to the explained methodology, in the period 1961-2020 20 dry years and 17 wet years were observed. The driest year was 2000, with z score -2.30 and total amount of precipitation of 367.7 mm. The wettest year was 2014, when 1095.1 mm of

precipitation was recorded (z=2.86). Three consecutive D years are observed at the beginning of the period (1961–1963) and in the last decade of the period (2011–2013). During the entire period, no more than two consecutive W years were recorded (1969–1970, 1977–1978, 1980–1981, 2004–2005, 2009–2010). From 1964-1968 a long sequence of N years was recorded. From 1998-2020, only five N years were recorded (Figure 6).

Table 7: Average amount of precipitation (mm) in Belgrade during two climate periods 1961-1990 и 1991-2020

Climatic Periods	Ι	II	III	IV	V	VI	VII	VIII	IX	Х	XI	XII	Annual
1961- 1990	49.3	44.4	49.5	58.8	70.7	90.4	66.5	51.2	51.4	40.3	54.3	57.5	684.4
1991- 2020	48.4	43.5	48.6	51.5	72.4	95.6	66.4	55.1	58.6	54.8	49.6	54.8	699.2

Source: Republic Hydrometeorogical Service f Serbia, Meteorogical yearbooks 1961–2020



Figure 6: Annual amount of precipitation and classification of years in Belgrade 1961–2020

Conclusions

In this research, temperature and precipitation data in Belgrade in the period 1961-2020 were analyzed. Presented results were obtained using two non-parametric tests: Mann-Kendall test and Pettit's test. The Mann-Kendall test with Sen's estimate indicates the existence of statistically significant trends and shows numerical changes on an annual basis. Pettit's test was used to identify changing points in the observed time series.

The results indicate an unequivocal increase of average annual and monthly temperatures. The increasing trend is statistically significant on annual level (0.042°C/year) and during all months except November. By applying the Pettit's test, we came to the result that

the changing point in the increase of the average annual temperature was 1997. The aforementioned trends can be explained by the effects of global warming, which have an impact on a regional and global scale.

The analyzed precipitation data for the same time series indicate that a trend without greater statistical significance is established. Statistically significant change at the annual level was not obtained. A statistically significant trends were detected in October (0.592 mm/year) and July (-0.416 mm/year). By monitoring the precipitation data in the research area, it was concluded that there is no significant changing point in the time series.

In order to reach a general conclusion about the change of climate elements, a comparative analysis of the data of two climate periods (1961–1990) and (1991–2020) was made. This indicates an increase in

both temperature and precipitation amounts in the second thirty-year period (1991-2020). The analysis of the inter-annual precipitation variability has shown that in the last two decades the number of N years has decreased and the frequency of D and W years has increased, which points to a possible increase of extreme climate events.

This research provides an insight into the dynamics of the climate in recent decades. It can be a firm base for the future study of climate changes in the area of Serbia, the Balkan Peninsula and Europe. Those research, firstly, must include the analysis of trends at the seasonal level, in order to accurately observe the time of year when greatest the trend decrease/increase occurs. In addition, a more complete analysis of the factors that influenced such a state of the climate should be done. Presented study should contribute a better understanding of recent climate change in Serbia. As recent studies have revealed (Tvardosz et.al., 2021), Europe continues to experience extremely cold winter months, which means that today's warming is a complex and spatially variable process (Hegerl et al., 2018; Krauskopf & Huth, 2020) that must be monitored on a continuous basis to assess its economic and social potential.

Funding

The study was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract number 451-03-68/2022-14/200091).

Conflicts of interest

The authors declare no conflict of interest.

Author contribution

Conceptualization, S.V. and M.D; methodology, U.D.; formal analysis, investigation and writing—original draft preparation, N.C; writing—review and editing, U.D., M.D, S.V. All authors have read and agreed to the published version of the manuscript.

References

- Ahmad I., Tang Deshan F., Wang T., Wang M. & Wagan, B. (2015) Precipitation trends over time using Mann-Kendall and Spearman's rho tests in Swat River basin, Pakistan. *Advances in Meteorology*. doi:10.1155/2015/43186
- Anđelković, G. (2007). Temperature conditions on July 2007 as extreme climatic phenomenon in Serbia. *Bulletin of the Serbian Geographical Society* 87 (2): 51-62.

https://doi.org/10.2298/GSGD0702051A [In Serbian: Анђелковић, Г. (2007). Температурне прилике у јулу 2007. године као екстремна климатска појава у Србији. Гласник Српског географског друштва 87 (2): 51-62. https://doi.org/10.2298/GSGD0702051A]

- Bajat, B., Blagojević, D., Kilibarda, M., Luković, J. & Tošić, I. (2014). Spatial analysis of the temperature trends in Serbia during the period 1961–2010. *Theor Appl Climatol* 121:289–301. DOI 10.1007/s00704-014-1243-7
- Bandyopadhyay, J., & Perveen, S. (2006). A scrutiny of the justifications for the proposed interlinking of rivers in India. *Interlinking of rivers in India: Overview and Ken-Betwa link*, 23.
- Branković, Č., Cindrić, K., Gajić-Čapka, M., Güttler, I., Pandžić, K., Patarčić, M., Srnec, L., Tomašević, I., Vučetić, V., & Zaninović, K. (2013). Sixth National Communication of the Republic of Croatia under the United Nation Framework Convention on the Climate Change (UNFCCC), Collection of papers,
- Burić, D., Luković, J., Ducić, V., Dragojlović, J., & Doderović, M. (2014a). Recent trends in daily temperature extremes over southern Montenegro (1951-2010), *Natural Hazards Earth Systems. Sci.*, 14, pp. 67-72.
- Croitoru A-E., Piticar A., & Burada, D-C. (2016) Changes in precipitation extremes in Romania. *Quaternary International* 415:325– 335.doi.org/10.1016/j.quaint.2015.07.028 1040-61
- Ćurić, V., Durlević, U., Ristić, N., Novković, I. & Čegar, N. (2022). GIS application in analisys of threat of forest fires and landslides in the Svrljiški Timok basin (Serbia). *Bulletin of the Serbian Geographical Society*, 102 (1), 107-130. https://doi.org/10.2298/GSGD2201107C
- Drazic, M. D., Veselinovic, M. M., Rakonjac, B. Lj.,
 Bojovic, R. S., Brasanac-Bosanac, B. Lj., Cule, M.
 N. & Mitrovic, Z. S. (2014). Geographic,
 landscape and the other natural characteristics of
 tourism. *European Journal of Geography Volume*5 (3): 96 -122. ISSN 1792-1341
- EEA report No 1 (2017) Climate change, impacts and vulnerability in Europe 2016. An indicatorbased report. European Environmental Agency http://www.eea.europa.eu/publications/climatechangeimpacts-and-vulnerability-2016
- Gajbhiye, S., Meshram, C., Singh, S. K., Srivastava, P. K. & Islam, T. (2015). Precipitation trend analysis of Sindh River basin, India, from 102-year record (1901–2002). *Atmospheric Science Letters* 17(1): 71–77. https://doi.org/10.1002/asl.602
- Gajić-Čapka, M., Cindrić, K., &Pasarić, Z. (2014). Trends in precipitation indices in Croatia, 1961– 2010. *Theoretical and Applied Climatology*, 117, 3-4, DOI 10.1007/s00704-014-1217-9
- Gavrilov, M. B., Marković, S. B., Janc, N., Nikolić, M., Valjarević, A., Komac, B., Zorn, M., Punišić, M., & Bačević, N. (2018). Assessing average annual air temperature trends using the Mann-Kendall test

in Kosovo. *Acta geographica Slovenica.* 58(1),pp. 8-25.

Gavrilov, M. B., Marković, S. B., Jarad, A., & Korać, V. M. (2015). The analysis of temperature trends in Vojvodina (Serbia) from 1949 to 2006. *Thermal Science* 19(2), pp. 339- 350.

Gavrilov, M. B., Tošić, I., Marković, S. B., Unkašević, M., & Petrović, P. (2016). The analysis of annual and seasonal temperature trends using the Mann-Kendall test in Vojvodina, Serbia. *Időjárás* 122(2), pp. 203-216.

Gocic, M., & Trajkovic, S. (2013). Analysis of changes in meteorological variables using Mann-Kendall and Sen's slope estimator statistical tests in Serbia. *Global and Planetary Change* 100: 172-182.

https://doi.org/10.1016/j.gloplacha.2012.10.014

Hegerl G-C., Brönnimann S., Schurer A., & Cowan T. (2018). The early 20th century warming: Anomalies, causes, and consequences. *WIREs Clim Change* 9. https://doi.org/10.1002/wcc.522

Helsel, D. R., & Hirsch, R. M. (2002). Statistical methods in water resources. In *Book 4, Hydrologic analysis and interpretation, Chapter A3* (Techniques of water-resources investigations of the United States Geological Survey). https://doi.org/10.3133/tm4A3

Hidmed (n.d.). Retrived from https://www.hidmet.gov.r-s/latin/meteorologija/stanica_moss1.php?moss_id=13274

IPCC, 2021: Summary for Policymakers. In: Climate Change 2021: The Physical Science Basis.
Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 3–32, doi:10.1017/9781009157896.001.

Jaiswal, R.K., Lohani, A.K. & Tiwari, H.L. (2015). Statistical Analysis for Change Detection and Trend Assessment in Climatological Parameters. *Environ. Process.* 2, 729–749.

https://doi.org/10.1007/s40710-015-0105-3 Kendall, M.G. (1962). *Rank Correlation Methods*.

Hafner Publishing Company, New York. Kendall, M.G. (1975). *Kendall Rank Correlation Methods*. Griffin: London, UK, 1-202.

Klapwijk, M., Csóka, G., Hirka, A., & Björkman, C. (2013). Forest insects and climate change: long-term trends in herbivore damage. *Ecol Evol* 3 (12):4183–4196. doi:10.1002/ece3.717

Krauskopf, T., & Huth, R. (2020). Temperature trends in Europe: Comparison of different data sources. *Theoretical and Applied Climatology*, 139, 1305–1316.

https://doi.org/10.1007/s00704-019-03038-w

- Kutiel, H., Luković, J. & Burić, D. (2015), Spatial and temporal variability of rain-spells characteristics in Serbia and Montenegro. *Int. J. Climatol.*, 35: 1611-1624. https://doi.org/10.1002/joc.4080
- Luković, B. J. (2013). Spatial pattern of climate change in Serbia in the instrumental period. Doctoral Dissertation. Belgrade: University of Belgrade, Faculty of Geography. [In Serbian: Луковић, Б. J. (2013). Увођење просторне димензије у проучавању колебања климе у Србији у инструменталном периоду. Докторска дисертација. Београд: Универзитет у Београду, Географски факултет, 1-204.]
- Luković, J., Bajat, B., Blagojević, D. & Kilibarda, M. (2014) Spatial pattern of recent rainfall trends in Serbia (1961–2009). *Reg Environ Change* 14, 1789–1799. https://doi.org/10.1007/s10113-013-0459-x
- Luterbacher, J., Werner, J.P., Smerdon, J.E., Fernandez-Donado, L., Gonzalez-Rouco, F.J., Barriopedro, D., Ljungqvist, F.C., Büntgen, U., Zorita, E. & Wagner, S. (2016) European summer temperatures since Roman times. *Environmental Research Letters*, 11, 1–12. DOI 10.1088/1748-9326/11/2/024001

Malinovic-Milicevic, S., Radovanovic, M. M., Stanojevic, G. & Milovanovic, B. (2016). Recent changes in Serbian climate extreme indices from 1961 to 2010. *Theoretical and Applied Climatology* 124: 1089-1098. https://doi.org/10.1007/s00704-015-1491-1

Mann, H. B. (1945). Nonparametric tests against trend. *Econometrica*, 13, 245-259.

Meteologos, (2012). Retrived from http://www.meteologos.rs/karakteristikegradske-klime-beograda/

Meteoblue (n.d.) Simulirani istorijski podaci o klimi i vremenu za Beograd. Retrived from). https://www.meteoblue.com/sr/vreme/historycli mate/climatemodelled/beograd_%D0%A1%D1% 80%D0%B1%D0%B8%D1%98%D0%B0_792680

Milanovic, M., Gocic, M. & Trajkovic, S. (2014). Analysis of Extreme Climatic Indices in the Area of Nis and Belgrade for the Period between 1974 and 2003. *Agriculture and Agricultural Science Procedia* 4: 408-415.

https://doi.org/10.1016/j.aaspro.2015.03.046

Milentijevic, N., Bacevic, N., Ristic, D., Valjarevic, A., Pantelic, M. & Kicovic, D. (2020). Application of Mann-Kendal (MK) test in trend analysis of air temperature and precipitation: Case of Macva district (Serbia). *Publication in Natural Sciences* 10 (1): 37-43.

https://doi.org/10.5937/univtho10-24774 NOAA National Centers for Environmental Information (2022), State of the Climate: Monthly Global Climate Report for Annual 2021.

https://www.ncei.noaa.gov/access/monitoring/m onthly-report/global/202113.

- Pešič, A. M., & Milovanović, B. (2016). Thermic regime and air temperature trends in Sumadija region (Serbia). *Journal of the Geographical Institute Jovan Cvijic SASA* 66(1):19-34. DOI: 10.2298/IJGI1601019M
- Pettitt, A.N. (1979). A Non-Parametric Approach to the Change-Point Problem. *Journal of the Royal Society: Series* C (Applied Statistics), 28, 126-135.
- Pohlert, T. (2015). Trend: Non-Parametric Trend Tests and Change-Point Detection, R package version 0.0.1. DOI: 10.13140/RG.2.1.2633.4243
- Popov, H., & Svetozarevich, J. (2021). Changes and contemporary trends in the annual amounts of precipitation in Serbia. *Journal of the Bulgarian Geographical Society*, Volume 44 (2021) 73–79. DOI 10.3897/j77102
- Radivojević, R. A., Martić Bursać, M. N., Gocić, J. M., Filipović, M. I., Pavlović, A. M., Radovanović, M. M., Stričević, S. Lj. & Punišić, R. M. (2015).
 Statistical analysis of temperature regime change on the example of Sokobanja basin in eastern Serbia. *Thermal science* 19 (2): 323-S330 https://doi.org/10.2298/TSCI150119019R
- Republic Hydrometeorological Service of Serbia RHMZ (2022). Meteorological yearbook climatological data 1961-2020.
- Salmi, T., Määttä, A., Anttila, P., Airola, T. & Amnell, T. (2002). Detecting trends of annual values of atmospheric pollutants by the Mann-Kendall test and Sen's slope estimates—the Excel template application MAKESENSE. *Helsinki: Finish Meteorological Institute*.
- Stojković, M., Plavšić, J. & Proharska, S. (2014).
 Long-term changes of annual and seasonal flows:
 A case of the Sava River. *Vodoprivreda*, 46, 39-48. [In Serbian: Stojković, M., Plavšić, J. & Proharska, S. (2014). Dugoročne promene

godišnjih i sezonskih proticaja: primer reke Save. *Vodoprivreda*, 46, 39-48.]

- Tošić, I., Hrnjak, I., Gavrilov, M. B., Unkašević, M., Marković, S. B., & Lukić, T. (2014a). Annual and seasonal variability of precipitation in Vojvodina. *Theoretical and Applied Climatology*, 117, pp. 331-341. doi:10.1007/s00704-013- 1007-9
- Trajkovic, S. & Kolakovic, S. (2009). Wind-adjusted Turc equation for estimating reference evapotranspiration at humid European locations. *Hydrology Research* 40 (1): 45–52. https://doi.org/10.2166/nh.2009.002b
- Trbić, G., Popov, T. & Gnjato, S. (2017). Analysis of air temperature trends in Bosnia and Herzegovina. *Geographica Pannonica*. 21 (2): 68-84 DOI: 10.18421/GP21.02-01
- Twardosz, R., Walanus, A. & Guzik, I. (2021). Warming in Europe: Recent Trends in Annual and Seasonal temperatures. *Pure Appl. Geophys.* 178, 4021–4032. https://doi.org/10.1007/s00024-021-02860-6
- Unkašević, M. (1994). The climate of Belgrade. Belgrade: Naučna knjiga. [In Serbian: Unkašević, M. (1994). Klima Beograda. Beograd: Naučna knjiga.]
- Vyshkvarkova, E., & Sukhonos, O. (2022). Compound Extremes of Air Temperature and Precipitation in Eastern Europe. *Climate*, 10, 133. https://doi.org/10.3390/cli10090133
- Westra, S., Alexander, L.V., & Zwiers, F.W. (2013). Global increasing trends in annual maximum daily precipitation. *J. Clim.*, 26 (11), 3904–3918. DOI: https://doi.org/10.1175/JCLI-D-12-00502.1
- Živanović, S., Ivanović, R., Nikolić, M., Đokić, M. & Tošić, I. (2020). Influence of air temperature and precipitation on the risk of forest fires in Serbia. *Meteorology and Atmospheric Physics* 132:869– 883. https://doi.org/10.1007/s00703-020-00725-6 http://www.meteologos.rs

https://climateknowledgeportal.worldbank.org https://www.meteoblue.com



Forum geografic. Studii și cercetări de geografie și protecția mediului
Volume XXII, Issue 1 (June 2023), pp. 16-27; DOI: 10.5775/fg.2023.066.i
© 2023 The Author(s). Published by Forum geografic.
Open Access article.
Creative Commons Attribution-NonCommercial-NoDerivatives License.

The changing river course and its impact on riverine society: A case study on the Padma River, Murshidabad District, West Bengal (India)

Dr. Sanatan GHOSH1*

¹ Associate Professor, Department of Geography, Krishnath College & Coordinator (Ad hoc) Department of Geography, Murshidabad University, Berhampore, Murshidabad, West Bengal, India.

* Corresponding author: sghosh2871972@gmail.com

Received on 20-01-2023, reviewed on 22-02-2023, accepted on 20-03-2023

Abstract

River bank erosion within Murshidabad district is a tragedy, which happens haphazardly along the bank of the Padma River system. The main objective of the study is to analyze the changing river course and its impact on society. By comparing the toposheet and Landsat (5 and 8) images of the Padma River of the Murshidabad district from the years 1924, 1990, and 2020, it is clearly observed that there have been significant changes on the southern portion of the river since 1924 (Bhagawangola II, Raninagar II, and Jalangi Block), while less changes occur on the central part of the river, which is near the Lalgola block. But in the year 2020 significant changes have been found in the northern and middle portion of the district near Samserganj and Lalgola block. Those parts belong to a highly vulnerable zone of bank erosion. This study analyzed some factors, such as soil stratification of the river bank and human encroachment as an obstruction to the natural river flow, also responsible for bank failure. The river has been consuming the vast portions of the right bank every year due to its increasing sinuosity. The victims are mostly in the Samserganj, Lalgola, Bhagawangola II, Raninagar II and Jalangi blocks, who suffered greatly from 1924 to 2020 due to moving of river course by 27% on the right side, which is notified from cross sectional analysis. Additionally, the predicted erosional impact of the region highlights the socio-economic perspective of the research area.

Keywords: dynamic river course, cross section, lateral erosion, Landsat, Padma River, socio-economic perspective

Introduction

Murshidabad district is situated in the moribund deltaic region of the lower Ganga plain in India. The deltaic rivers have the tendency to oscillate within a wide limit. The alteration of river channels is almost a universal occurrence. Murshidabad district's people have probably been affected by the river for a long time. Murshidabad was the capital and important Business centre of Bengal, Bihar and Orissa in the British period. Murshidabad has lost its earlier glory due to historical and geographical reasons. Another historic town that perished naturally was Karnasubarna. The probable cause may be the severe erosion of the Bhagirathi River or the lack of a sufficient water supply when the main flow of the Ganga was diverted through Padma. Researchers have been studying the geomorphological phenomenon of river bank erosion and channel shifting for the past few years (Mukhopadhyay & Das Gupta 2010; Parua, 2006, 2009, 2010; Rudra, 2005, 2010; Ghosh 2015; Islam et al., 2019; Rahman, 2010; Raman et al., 1986). The changing river courses also led to river bank erosion, devastating floods, extensive silt deposition along river beds, meandering cut-off which are the serious threat to human habitation as well as

environment. The river system is made up of mainly the Ganges, the Padma, the Bhagirathi, Jalangi, and Bhairab. These river systems are constantly altering their paths, and deltaic rivers do so in a wide range. The river systems of the district are generally meandering and braided in nature. Rivers help the systems in dynamic equilibrium. When river channels are altered due to naturally dynamic hydrologic conditions, the river readjusts itself in terms of size, profile, and pattern to re-establish its previous balance or equilibrium (Couture, 2008). The free-flowing river is always adjusting the equilibrium condition through erosion, transportation, and deposition processes. When the deposition process is underway on one side, the erosion process is underway on the other bank side. Mid channel bar or char formation is restricted to mid channel flow pattern and flow concentrated to bank side, as result of severe bank erosion. The erosion and deposition of a meandering river are continuously characterized by the processes of emergence, submergence, and reemergence. This phenomenon has been explained as follows: "accumulated silt leads to the rise of a sand-bed in the interfluves (the region of higher land between channels); being obstructed by this bed of the river then divides into two channels with the sand-bed in between. This makes the flow obligue. The flow gets obstructed