

Forum Geografic - Studii și cercetări de geografie și protecția mediului (FG - S.C.G.P.M.)

The journal *Forum geografic. Studii de geografie și protecția mediului* (Geographical Forum – Geographical Studies and Environment Protection Research) was founded in 2002, and it seeks to publish high quality research in the domains of geography, environment protection and other related domains. It addresses a range of issues, such as geomorphology, pedology, climatology, hydrology, human geography and environment. Its content is directed to a broad audience, including both academics and policymakers. The

papers selected for publication in the journal are subject to a review process using references from universities worldwide. The journal is currently indexed by the following databases: SCOPUS, DOAJ, EBSCO, ERIH PLUS, Index Copernicus, Scipio, CrossRef, Ulrichsweb, IGU, Google Scholar, WorldCat, ProQuest, ZDB, CNCSIS, DRJI, citefactor.org journals indexing, InfoBase index, Eurasian Scientific Journal Index, ResearchBib.

Editor-in-chief:

Sandu BOENGIU, Physical Geography, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Liliana POPESCU, Human Geography, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Executive editor: Oana Mititelu Ionuș, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Technical editor: Cristiana Vlcea, Geography Department, University of Craiova, 13, Al. I. Cuza Street, Craiova, Romania

Editorial Advisory Board:

Iuliana ARMAȘ, *University of Bucharest, Romania*

Sorin AVRAM, *University of Craiova, Romania*

Lucian BADEA, *The Institute of Geography, The Romanian Academy*

Dan BĂLTEANU, *The Institute of Geography, The Romanian Academy*

Zeljko BJELJAC *Geographical Institute Jovan Cvijić, Serbia*

Lucian BLAGA, *University of Oradea, Romania*

Sandu BOENGIU, *University of Craiova, Romania*

Léon BRENIG, *University of Brussels, Belgium*

Manola BRUNET, *University of Tarragona, Spain*

Sorin CHEVAL, *Henri Coandă Air Force Academy, Brașov, Romania*

Laura COMĂNESCU, *University of Bucharest, Romania*

Lóczy DÉNES, *University of Pécs, Hungary*

Philippe DONDON, *Ecole Nationale Supérieure E.I.T.M.M. de Bordeaux/ Institut Polytechnique de Bordeaux, France*

Monica DUMUTRAȘCU, *The Institute of Geography, The Romanian Academy*

Dan DUMITRIU, *Al.I.Cuza University, Iași, Romania*

Recep EFE, *Balikesir University, Turkey*

Robert FOVELL, *University of California, USA*

Teodoro GEORGIAIDIS, *Institute of Biometeorology (IBIMET), Bologna, Italy*

Adrian GROZAVU, *Al.I.Cuza University, Iași, Romania*

Nelly HRISTOVA, *St. Kliment Ohridsky University of Sofia, Bulgaria*

Ioan IANOȘ, *University of Bucharest, Romania*

Alexandru ILIEȘ, *University of Oradea, Romania*

Mihaela LICURICI, *University of Craiova, Romania*

Gabriel MINEA, *Institutul Național de Hidrologie și Gospodărire a Apelor, Romania*

Oana MITITELU IONUȘ, *University of Craiova, Romania*

Emil MARINESCU, *University of Craiova, Romania*

Mirela MAZILU, *University of Craiova, Romania*

Ciprian MĂRGĂRINT, *Al.I.Cuza University, Iași, Romania*

Alexandru NEDELEA, *University of Bucharest, Romania*

Amalia NIȚĂ, *University of Craiova, Romania*

Zvi Yehoshua OFFER, *Ben-Gurion University, Israel*

Maria PĂTROESCU, *University of Bucharest, Romania*

Aurel PERȘOIU, *Emil Racovita Institute of Speleology, Romanian Academy*

Dănuț PETREA, *University of Babeș-Bolyai, Cluj-Napoca, Romania*

Liliana POPESCU, *University of Craiova, Romania*

Kumar Rai PRAVEEN, *Banaras Hindu University, India*

Maria RĂDOANE, *Ștefan cel Mare University, Romania*

Milan RADOVANOVIĆ, *Geographical Institute Jovan Cvijić, Serbia*

Khairav RAVINDRA, *School of Public Health, PGIMER, Chandigarh, India*

Daniela ROȘCA, *University of Craiova, Romania*

Daniel SIMULESCU, *Agentia Nationala pentru Arii Naturale Protejate*

Igor SIRODOEV, *Universitatea "Ovidius" din Constanța, Romania*

Fábián Ákos SZABOLCS, *University of Pécs, Hungary*

Cristina ȘOȘEA, *University of Craiova, Romania*

Magdy TORAB, *Alexandria University, Egypt*

Marcel TÖRÖK – OANCE, *West University of Timișoara, Romania*

Cristiana VÎLCEA, *University of Craiova, Romania*

Alina VLĂDUȚ, *University of Craiova, Romania*

Nenad ŽIVKOVIĆ, *Belgrade University, Serbia*

Martina ZELENKOVA, *Technical University of Kosice, Slovakia*

Zbigniew ZWOLIŃSKI, *Adam Mickiewicz University (Poznan), Poland*

Associate Editors:

Slavoljub DRAGIĆEVIĆ, Faculty of Geography, Belgrade University, Studentski trg 3/3, Belgrade, Serbia

Vesna LUKIĆ, Demographic Research Centre, Institute of Social Sciences, Kraljice Natalije 45, Belgrade, Serbia

Nina NIKOLOVA, Faculty of Geology and Geography, "St. Kliment Ohridsky" University of Sofia, Tzar Osvoboditel Blvd. 15, Sofia, Bulgaria

Founding Editors: Boengiu S., Marinescu E., Pleniceanu V., Tomescu V., Enache C.

Cover photo: *Făgăraș Mountains, Sibiu county (by Alina Vlăduț)*

For instructions for authors, subscription and all other information please visit our website

<http://forumgeografic.ro>

before submitting any papers please select the section *Publishing rules* from the About page and read thoroughly the submission instructions for authors

ISSN 1583-1523 (print)
ISSN 2067-4635 (online)
DOI prefix: 10.5775

Instructions for Authors

Article submission

In order to disseminate the research results in the field, researchers, scholars and professionals are welcome to submit an electronic version of the manuscript (in Microsoft Office Word format) to the editorial office (forum.geografic@gmail.com).

Submission requirements: The submission of an article for publication in our journal implies that the article has not been published before, nor it is being considered for publication in other journals. Authors are responsible for the content and the originality of their contributions. In order to be published, articles must be thoroughly researched and referenced.

IMPORTANT: All papers must be submitted in electronic format, only in English language.

Copyright statement

By submitting a scientific work to *Forum geografic* the submitters agree to declare the following:

- the submitted work belongs exclusively to the declared authors;
- the submitted work represents original scientific research;
- the submitted work has not been published or submitted for publishing to another journal;
- if the submitted work is published or selected for publishing in *Forum geografic*, the authors waive any patrimonial claims derived from their authorship for the submitted work; the authors retain the moral rights for their submitted work, as granted under the Romanian applicable law; also, the authors agree to refrain from ulterior submitting of the work to other journals.

The submitters agree to be solely held accountable in case of breaching the above terms and to defend the representatives of *Forum geografic* in the event of a lawsuit related to the submitted work.

When submitting a paper the authors are required to print, fill and send a scanned copy of this declaration.

Privacy statement

The submitted personal data, such as names or email addresses, are used only for the declared purpose of the *Forum geografic* journal (publishing original scientific research) and are not available to third parties.

Manuscripts are received at all times. However, in order to have your article published in the current year, the manuscripts must be submitted until the 15th of February for the first issue of the current year and until the 1st of September for the second issue.

Article format

All manuscripts must be edited entirely in English. Articles must include:

- Title
- Author's name(s). For each author you must mention the author's scientific title, his affiliation (institution) and e-mail address;

- Abstract (maximum 300 words);
- Keywords (not more than 5-6 words);
- Acknowledgments (if any);
- Main body of text (structured according to Introduction, Data & Methods, Results & Discussions, Conclusions);
- Illustrations (graphs, diagrams, maps, photos – should have indications of their positions in the text and title written in English) must be also submitted in electronic format, preferably in JPG, PNG or BMP format and must be referred to as Figures, which should be numbered with Arabic numbers.
- Tables must be numbered with Arabic numbers and should not repeat data available elsewhere in the text.
- References must be indicated in the text, between brackets and they must include the author's name and the date of the publication (Popescu, 2000). When three or more authors are referred, they will appear in the text as follows: (Popescu et al., 1997). References must be listed in alphabetical order at the end of the text.

The following style sheet is recommended:

- for journals:

Miletić, R., Lukić, V., & Miljanović, D. (2011). Deindustrialization and structural changes in commuting flows in Serbia. *Forum geografic*, X(2), 244-254. doi:10.5775/fg.2067-4635.2011.009.d

- for books:

Bran, F., Marin, D., & Simion, T. (1997). Turismul rural. Modelul european, Editura Economică, București

- for papers from conference proceedings:

Deci, E. L., Ryan, R. M., (1991), A motivational approach to self: Integration in personality. In R. Dienstbier (Ed.), *Nebraska Symposium on Motivation: Vol. 38. Perspectives on motivations* (pp. 237-288). Lincoln: University of Nebraska Press.

Review process

All the manuscripts received by the editors undergo an anonymous peer review process, necessary for assessing the quality of scientific information, the relevance to the field, the appropriateness of scientific writing style, the compliance with the style and technical requirements of our journal, etc. The referees are selected from the national and international members of the editorial and scientific board, as well as from other scholarly or professional experts in the field. The referees assess the article drafts, commenting and making recommendations. This process leads either to acceptance, recommendation for revision, or rejection of the assessed article. Editors reserve the right to make minor editorial changes to the submitted articles, including changes to grammar, punctuation and spelling, as well as article format, but no major alterations will be carried out without the author's approval. Before being published, the author is sent the proof of the manuscript adjusted by editors. If major revisions are necessary, articles are returned to the author so that he should make the proper changes. Authors are notified by email about the status of the submitted.

Risk of Dynamics of the River Stream in Tectonic Areas. Case studies: Curvature Carpathian - Romania and Maghreb Chain - Algeria

Florina GRECU^{1,*}, Chaouki BENABBAS², Mădălina TEODOR³, Manel YAKHLEFOUNE²,
 Iulian SĂNDULACHE¹, Nabil MANCHAR⁴, Takki Eddine KHARCHI², Gheorghe VIȘAN¹

¹ University of Bucharest, Faculty of Geography, Bucharest, Romania

² Constantine University 3, Institute of Urban Techniques Management, Constantine, Algeria

³ University of Bucharest, Faculty of Geography/Romanian Young Academy (RYA), Bucharest, Romania

⁴ University of Oum El Bouaghi, Department of Geology, Constantine, Algeria

* Corresponding author: florinagrecu@yahoo.com

Received on 19-12-2020, reviewed on 02-02-2021, accepted on 05-05-2021

Abstract

The tectonic characteristics of the Alpine orogeny highlight some similarities in the genesis of landforms over large areas. This paper analyses the role of tectonic processes in two subsiding areas with uplift movements. The overall tectonic activity implies the formation and evolution of the hydrographic network and its correlation with the major relief structure, thus forming streams parallel to the high-altitude lines, as well as consistent, transversal, and subsequent streams.

These tectonic processes are accompanied by antecedents and/or epigenesis common in the Eastern Carpathians, Romania. Hence the Bistrița River is longitudinal compared to the Bistrița and the Stânișoara Mountains, but also transversal in other sectors. In Algeria, the Northeast, especially the Zighoud Youcef zone (NE of Constantine), is characterized by a particular morphography and morphometry of the river basins which reveals reorganizations of the river network through their transversal or longitudinal characteristics compared to the structure and tectonics of the region. In areas with imminent catchments, the regressive erosion amplifies the slope dynamics and slope processes vulnerability.

The uplift and subsidence movements impose reorganizations of the hydrographic network of the catchments with the alternation of the erosion and sedimentation processes it creates a certain fluvial style pattern. In the subsidence of the Întorsura Buzăului depression, the longitudinal profile of the Buzău River has a very low slope and its meanders create islets. This is subject to frequent floods which imposed the riverbed regularization. However, in Zighoud Youcef, the Oued Smendou River also indicates some similarities with the Buzău River structure due to subsiding areas of the Mila-Constantine basin.

Keywords: *tectonic risk, Alpine orogeny, river network, Constantine region, the Atlas, The Curvature Carpathians*

Rezumat. Riscul dinamicii cursurilor râurilor în ariile tectonice. Studii de caz: Carpații de Curbură (România) și lanțul Maghrebian (Algeria)

Caracteristicile tectonice ale orogenului alpin permit evidențierea unor similitudini în geneza formelor de relief pe spații extinse. Lucrarea analizează rolul manifestărilor tectonice în două arii subsidente și tectonic pozitive. Tectonica de ansamblu impune formarea și evoluția rețelei hidrografice în raport cu structura majoră a reliefului; astfel, se formează: cursuri paralele/longitudinale cu linia marilor înălțimi și cursuri transversale/subsecvente. Procesele sunt însoțite de antecedente și/sau epigeneze, frecvente în Carpații Orientali. Astfel, râul Bistrița este longitudinal fata de Munții Bistriței și Munții Stânișoara, dar și transversal în alte sectoare. În Algeria, Atlasul de nord-est, în special zona Zighoud Youcef (NE de Constantin), se caracterizează printr-o morfografie și morfometrie particulară a bazinelor hidrografice, care dezvăluie reorganizările rețelei fluviale prin caracteristicile lor transversale sau longitudinale în comparație cu structura și tectonica regiunii. În zonele cu captări iminente, eroziunea regresivă amplifică dinamica pantei și vulnerabilitatea proceselor de pantă.

Mișcările de înălțare și mișcările subsidente determină reorganizări ale rețelei hidrografice prin captări, alternanța proceselor de eroziune și acumulare și deci crearea unui anumit tipar fluvial. În depresiunea Întorsura Buzăului, profilul longitudinal al râului colector are o pantă foarte mică, favorizând procesele de meandrare și crearea ostrovelor. Ca urmare a riscului inundațiilor, s-au întreprins ample lucrări de amenajare a albiei minore. În arealul Zighoud Youcef, râul Oued Smendou indică, de asemenea, unele asemănări cu structura râului Buzău datorită zonelor de subsidență din bazinul geologic Mila-Constantine.

Cuvinte-cheie: *risc tectonic, orogeneza alpină, rețea hidrografică, regiunea Constantin, Munții Atlas, Carpații de Curbură*

Introduction

The vulnerability to the active tectonic processes indicates a tectonic risk with environmental and societal consequences. These risks are usually analysed as geomorphological, geological and hydrological, according to the environment and the main exogenous agents (relief/gravity, water/floods, soil/erosion, etc.). In order to study those risks, it is

very important to know the geological, tectonic and paleo-evolutionary characteristics, including the paleoclimatic factors of the study area to predict the triggering factor and the dynamics of the extreme hazardous processes. This paper analyses the principal role of the tectonic processes significant for triggering a risk in both areas of interest of the Alpine orogeny, the Carpathian and the Mila-Constantine basin. Alpine orogenic regions require comparative

studies due to tectonics and structure reflected by the arrangements of peaks influenced by active uplifting movements in addition to the presence of depressions with subsiding movements (Ciccacci et al., 1999; Benabbas, 2006).

The overall tectonics involves the formation and evolution of the hydrographic network in relation to the major structure of the relief, against the general background of the Quaternary climate fluctuations (Piacentini and Miccadei, 2014; Piacentini et al., 2015). Thus, parallel watercourses to the line of great heights (consistent) and transversal watercourses (subsequent) are formed. In general, the transversal watercourses formed by antecedents are well highlighted on the physical map of Romania (Fig. 1) (Naum 1961, Grigore 1989, Orghidan 1939, Posea 2005).

The processes are followed by antecedents and/or epigenesis, common in the Eastern Carpathians. Thus, the Bistrița River is longitudinal compared to the Bistrița and Stânișoara mountains while transversal in other sections (Donișă 1968, Săndulache 2007) (Fig. 2 B).

In the north-east of Algeria, it is necessary to take into consideration the heritage and the geological history to explain the formation and the evolution of the relief, thus, the Rhumel and the Ben Brahim-Bouhadjeb and Smendou rivers are longitudinal compared to the Numidic chain while transversal in other sections.

However, the north of Algeria is characterized by a varied and complex geological and geomorphological context, distinct by a strong tectonic and an active seismic activity (Benabbas, 2006; Yelles-Chaouach et al., 2006; Benzid, 2017). The Mila-Constantine basin in north-east of Algeria, is more susceptible to landslides and floods hazards than the other parts of the country which is presented by the record of many important events (Bourenane et al., 2014, 2016; Manchar et al., 2018 Manchar et al 2019). These phenomena become more complex where the impact of the hydrogeomorphic hazard evolve in space and time (Hamadou and Amireche, 2018).

Based on field research and bibliographic documentation, applying multicriteria geomorphological methods as well as GIS techniques, this paper highlights two cases: 1) the dynamics of a representative river from the Curvature Carpathians – the Buzau River. 2) the characteristics of the drainage network related to the morpho structure and tectonics in the north-eastern part of Algeria (Constantine region). The analyses of the vulnerability of rivers to extreme natural processes, initially due to tectonic processes (Benabbas et al., 2017). Morphotectonic and morphostructural risk is the result of active tectonics that creates a typical landscape that attests to the degree of danger

(Dramis F, Cliff Oliver 2016) However, this argument is particularly useful in establishing the long-term effect of anthropogenic actions in the slope and river (dams, bridges, skiing... etc.) (Teodor 2013; 2017). The alpine regions create a topographic contrast between mountain peaks and depression basins even in mountain regions of medium altitudes (Raffy Jeannine 1983; Peulvast J-P et al. 2013).

The dynamics of the rivers is seen in a hydro-geomorphological context, of the synergistic relations between the river/water and the topographic/relief surface (Ballais et al., 2011, Grecu et al., 2017b; Arseni et al., 2019, Sembroni et al. 2021). Precipitation and changing climatic conditions increase runoff variability and control its evolution, which have a major impact on hydro-geomorphological hazard. That, in turn, may explain the intensity of some tectonics and flood events (Meddi et al., 2009; Waleed A 2021).

Study area – vulnerability factors

Romanian Carpathians

The Romanian Carpathians, or South-Eastern Carpathians (according to Mihăilescu, 1963), are the highest Alpine mountains in Europe (about 1500 km between the Vienna Pass and the Timok River) (Institute of Geography, Geography of Romania 1987). On the Romanian territory, the orogenic, geological and morphological elements determined the separation of the subunits (Fig. 1A and 1B).

The Eastern Carpathians (including the Curvature Carpathians) with a maximum altitude of 2303 meters in Pietrosul Rodnei peak, characterized by approximately parallel peaks, volcanic (Neogene volcanism in the west (Fielitz and Seghedi 2005), crystalline (center) and sedimentary (East); the Southern Carpathians (mainly crystalline schists, limestones and conglomerates towards the limits of the massif, with a maximum altitude of 2545 meters in the Moldoveanu peak and the Western Carpathians (maximum altitude of 1849 meters in Cucurbăta Mare peak, Biharia Massif) characterized by the presence of old volcanism, crystalline in high massifs (Mesozoic sedimentary). The name of the mountain sectors is considered by the position within Romania, but also towards the Transylvanian sedimentation basin, from the central part, a hilly plateau with slight subsidence movements. The hydrographic network was formed in different stages, taking into account the arrangement of mountain peaks, the ratios to baseline levels, climatic conditions (Posea, 2005), also depending on the capture, epigenetics, antecedents, erosional and cumulative have finalized the morphology of the rivers and implicitly of the hydrographic basins, so that today it presents a quasi-circular aspect compared to the Romanian

Carpathians system (Posea et al., 1974). Some of the most complex genetic aspects are presented by the rivers of the Eastern Carpathians (Fig. 2A).

The Romanian Carpathians are an arcuate chain of mountains, formed in response to the continental collision between the European Plate and several microplates during the Alpine Orogeny (Schmidt et al., 1998; Mațenco and Bertotti 2000, Tarapoanca et al. 2003). The main depocenter in the Carpathians foredeep is the Focșani Depression, where subsidence continues up to present time. During the Pliocene–Quaternary, the western flank of the Siret subsidence region (Focșani depocenter) was tilted as a

consequence of the uplift of the neighbouring Carpathian chain (Necea et al., 2005; Mațenco et al., 2007; Necea 2021). Its opposite important regional vertical movements: the uplifting chain and subsiding foredeep (about +/-5 millimetres per year; (Zugrăvescu et al., 1998) at less than 100 km in distance (Fig. 3). It is also characterized by the outcrop of rock types susceptible to erosion: sand, clay, conglomerate and marls that record the change from a marine to fluvial-lacustrine environment (Molin et al., 2012; Teodor et al., 2014; Teodor and Dobre, 2015).




Fig. 1: A) Geographical position within Romania and large geographical units; B) Upper Buzău and Întorsura Buzăului Depression - physical-geographical map




Fig. 2: Geological maps of Romania. Carpathian orogeny units and the hydrographic network in the central part of Romania (A) and geological structure (B); the longitudinal and transversal character of the rivers in the Eastern Carpathians - the Bistrița basin

The complex and orderly structure, in rows arranged on the north-south direction (from east to west – sedimentary, crystalline, volcanic), separated by large depressions, is the main factor in deciphering the hydrographic network, in which the longitudinal sections alternate with the transversal ones. The southern part of the Eastern Carpathians presents an arch of the sedimentary ridge (the crystalline and volcanic series are missing) with its own morphological structure, imposed by the Black Sea microplate (Posea, 2005).

In the Curvature Carpathians, the recent vertical crustal movements (elevation-subsidence at short distances), the existence of the most active epicentral area for earthquakes in Romania and continental Europe, the relief modelling and the evolution of the hydrographic network are made according to their own patterns (Grecu, 2016) (Fig. 3A).

The morphological complexity of the Curvature Carpathians also imposes itself on the external marginal relief (Grecu, 2010). The contact area between the Sub-Carpathian and the Romanian Plain is represented by a piedmont belt (glacis-like, macro-

landform associated with the monocline), traditionally considered a part of the Romanian Plain because of its very gentle slopes. The subsidence of the Romanian Plain coupled with the Carpathian and Sub-Carpathian uplift induced a regional base level, lowering that allowed the generation of fluvial terraces and entrenched alluvial fans interacting with the Quaternary climate changes (Grecu and Sacrieru 2009; Grecu et al., 2007, 2010; Grecu et al., 2014). The element of complexity is also due to local tectonic features (Necea, 2021), like the anticlines relative to the so-called "Wallachian" phase (Săndulescu, 1988).

These regional features have an influence on rivers crossing the Curvature area. Several rivers of this area have concave and steep longitudinal profiles and they are in a transient state of disequilibrium as a consequence of a more recent emersion of the Curvature Carpathians (Badea and Niculescu, 1964; Molin et al., 2012). Moreover, other previous studies showed that these rivers have the highest suspended sediments loads in Romania (up to 25 tons per hectare per year) (Mociorniță and Birtu, 1987 cited by Zaharia et al., 2011; Dumitriu, 2020).





Fig. 3: A) Recent vertical crustal movements map of South-Eastern Romania (after Zugrăvescu et al. 1998). Legend: 1 – seismic lines (mm/year); 2 – active crustal faults; 3 – areas affected by diapirism; B) Curvature Carpathians – geological map (data processed from the 1: 200,000 vector geological map)

Buzău River – river formation, vulnerability factor

The Buzău River springs from the Carpathian flysch, the northern mountain side of the Ciucas Mountains, from an altitude of approximately 1800 meters, having a south-north direction. From the source area, to the confluence with Acris (about 20 km), the river follows a geological contact between the less resistant and conglomerate Cretaceous formations with a slope of over 60 m/km to the confluence with Strâmbu, 14 m/km at Vama Buzău (State Geology Committee. Geological map, Covasna

sheet 1970). From Acris the river is oriented to the northeast enters the tectonic-erosive depression, where the slope of the river is reduced, about 4-4.5 m/km the accumulation processes lead to the formation of islets and frequent floods. Approximately from the city centre, it is oriented to the South-East and then to the South (with local modifications). The river basin, covering an area of 5264 km², develops within the Curvature Carpathians and the Curvature Sub-Carpathian, and within the Romanian Plain. It flows into the Siret River with a length of 334.4 km (Uivari, 1972). The longitudinal profile presents sections, with varying intensity of the relations

between erosion and accumulation (Fig. 12) influenced by the relations with the slopes (in the plain at Bănița at 0.7 m/km) (Minea, 2011).

Within the mountain area of the Buzău River, there are differentiated five sectors (Grecu et al., 2017 a; Grecu et al., 2018):

1. Întorsura Buzăului sector (between Buzău-Afiniș junction, (Brădet settlement) and Sita Buzăului village) is distinguished by the active dynamics of the river bed and by hydrotechnical works against floods, regularization of the minor river bed and shore defence (Fig. 4A and 4B);

2. Northern Transcarpathian section (between Sita Buzăului and the Buzău-Harțagu junction (partly Buzău Gorge, corresponding to the line of great heights);

3. The internal southern Transcarpathian sector (Harțagu/Băile Siriu – Siriu village, including the Rockfill Dam);

4. External Transcarpathian – Sub Carpathian sector located between Siriu village and downstream of Pălțineni, at the entrance into the Sub-Carpathians.




Fig. 4: A) Minor River bed with river deposits at Brădet (Întorsura Buzăului); B) The transverse valley of the lip in the Harțagu gorges

The Buzău River was formed during successive stages, with specific morphologies, in correlation with the geological and tectonic aspects of the Curvature Carpathians. The most discussed stage is related to the bend formed by the Buzău at Întorsura Buzăului and therefore the change of flow direction towards it (Orghidan, 1969; Posea and Garbacea, 1959; Posea, 1971, 2005).

Here, we address the following issues:

- 1) The problem of the transversal course;
- 2) Formation of the course between spring and Bradet (confluence with Acris);
- 3) The course of the river within the Întorsura Buzăului Depression.

The rivers from the Curvature Carpathians are transversal rivers, some bypass the line of the great heights (see the geological map and the physical map). The Buzău River forms in the line of the great heights (Siriu-Penteleu) a narrow valley type key between the confluences with Crasna and Harțagu. In fact, the morphological characteristics, the alternation of basins with narrow sectors, the analysis of the longitudinal profile and the analysis of the levels (Nordon, 1933) are arguments in the antecedent formation of the Transcarpathian Buzău (Orghidan, 1969; Posea, Garbacea, 1959; Posea 1971, 2005). This aspect is also reflected in the dynamics of the slopes, the activation and/or reactivation of the slope processes (landslides, erosion).

The upper lift and the watercourse of the Întorsura Depression are influenced by the tectonic movements – subsidence in the bottom of the depression, of the rise in the southern part on the line of the great heights and by the geological structure (Orghidan, 1939; Naum, 1961; Posea, 1971, 2005; Ielenicz, 1984; Grigore, 1989). Upper Buzău, Bâsca Mare and Bâsca Mica developed their basins over the current line of high altitudes, the evidence being the erosion surface of +/-1000 meters. According to Orghidan (1969 p. 43), there are no traces to prove that Buzău river is a late intruder.

The organization of the drainage network was done differently as claimed by some authors, and their opinions must be taken into account when considering the different stages in the evolution of the river. Thus, the influence that the lower base level of the Brașov Depression had on the disorganization of a paleo-network is accepted. The first stage refers to the existence of a paleo-Buzău with springs in Harghita, over the Brașov Depression (Posea and Garbacea, 1959), and explains a paleo network with the right tributary of the Upper Buzău in Ciucaș. The subsidence in the Turn Depression determined the concentration of the hydrographic network towards a local level, and then the reorganization according to the game of the basic levels and of the Quaternary tectonic movements (Iancu, 1971; Ielenicz, 1984). In our opinion (Grecu), the oscillations of the riverbed

on a river-lake subsident relief imposed a gradual direction of the flow to the south on easy lines. The Buzău Valley corridor is located on an axial sinking area of the folds, it is a tectonic corridor (Socolescu et al., 1975).

Along the corridor, there is an alternation of narrow transverse and diagonal sections, created in the Miocene and the Pliocene and also elongated depressions corresponding to synclines. This orientation is still observed in the dynamics of the riverbed. The analysis of meanders (1980 and 2017) from the source to the confluence with Crasna shows changes by sectors as follows (Grecu et al 2021):

- In the upper watercourse (up to Brădet), the meandering index increases from 1.35 in 1980 to 1.4 in 2017);

- In the depression area (between Brădet and Sita Buzăului), the lateral erosion is the most active, the meanders with variable lengths are bigger and more numerous, some of them have not been maintained; the meandering index increases in 1980 to 1.46; in 2017 it remains the same, but decreases compared to 1980 to 1.4;

- Downstream, towards the confluence with Crasna, the meanders are dense but of much smaller lengths (coefficient in 1980 and in 2017 of 1.26).

Overall, from the source to the confluence with Crasna, the meandering index decreases, but it has higher values in the central part of the subsidence area. The transverse profiles highlight the leftward movement of the active area.

Thus, for 1980, the largest meander displacement is 2263.02 meters to the left in the upper sector I (up to Brădet), 1079.9 meters to the left in sector II depression and 667.70 meters to the left in sector III Zăbrățu-Crasna (upstream of the gorge). For 2017, the largest shift of meanders compared to the asymmetry of the river is also to the left and is 2161.59 meters in sector I, 1078.81 meters in sector II and 667.66 meters in sector III, but there are slight decreases in values due to regularization of watercourses. Higher displacement values can be attributed to local petrographic conditions, in conditions of a lower sinuosity index.

Geomorphological and geological settings of Mila-Constantine Neogene basin

The area concerned by this study is located within the Mila-Constantine Neogene basin. It is part of a group of intramountain basins that characterize the Alpine chain of North-East of Constantine which was mainly structured during the Tertiary phases as a result from the structuring of the Maghrebids Basin and its African and European margins (Benabbas, 2006). It is a post-nappe basin that has been filled by a silico-clastic and carbonate series of Mio-plio-Quaternary age (Vila, 1980; Coiffait, 1992). This

receptacle is installed on a substratum pre-structured by thrust sheet units of the Tellian, Penitellian and neritic units belonging to the external zones of the Maghreb chain of eastern Algeria (Vila, 1980; Wildi, 1983).

Our study area is located on several distinct paleogeographic and structural domains: internal domain, Flysch domain, and external domain. The Maghreb chain is an Alpine segment characterized by the accumulation of large thrust sheets with south vergence (Benabbas, 2006).

The altitude ranges from 500 to 1200 meters above sea level. This area is characterized by a dense hydrographic network constituted by a set of oueds (rivers), ravines and gullies. This network is mainly represented by the following oueds: Oued Smendou located in the Kébir Rhumel-Smendou watershed, which is the principal stream in the basin (Fig. 5), and Oued Benbrahim-Bouhadjeb located in the Safsaf watershed (Fig. 5), and Oued El Kram, and Oued El Aria in the east which has permanent flow. Often, these streams are flowing to the Northeast direction.

The study area is characterized by a semi-arid climate with high temperatures (from 28 to 41°C) and low precipitations (from 600 to 900mm) in the summer, and high humidity, precipitation and low temperature in the winter. We can speak of two typical seasons (rainy and dry) that stand in contrast. About 63% of the annual rainfall quantity is concentrated between December and February each year. (Benzid, 2017; Manchar, 2021)

At the orographic level, the study area can be subdivided into three distinct natural sub-areas: the first one is the mountainous massifs of the Numidian chain in the north, the second one is the hills of Conde Smendou and Zardezas with altitudes varying from 400 to 500 meters and the third one is the low areas which are represented by:

1. The depression of Oued Beni Brahim-Bou Hadjeb, which drains a large part of the southern part of our study area and seems to follow a West-East direction at the beginning to change to a South West - North East, and finally to reach the valley of Oued Safsaf (beyond the mountains of Ouled Hababa) (Fig. 5, 7);

2. The depression of Oued Smendou which in its upstream part Oued Smendou is presented as a geomorphological, hydrographic, and geological (tectonic anomaly), indeed, this Oued was supposed to converge on Oued Beni-Brahim – Bou Hadjeb to join Oued Safsaf, however, it suddenly changes direction to the south of the town of Zighout Youcef forming a perfect angularity and takes a north-east south-west direction to join Oued Rhumel-Kébir (Fig. 8, 9). (Benzid, 2017; Benabbas et al., 2019; Benabbas et al., 2021).

The slopes that constitute the Mio-plio-Quaternary filling, are juxtaposed on several kilometres and are

composed of formations with clay predominance (Raoult, 1974).

From the morphological point of view, the basin of Mila-Constantine is divided, from west to east, in three units – Ouled Kebbeb, the Smendou unit that is analysed in this work, and the Mila unit, which is the most important. These units are inserted between the Numidic chain in the north and the mountains of Constantine in the south (Fig. 9).

Geologically, the study area is characterized by superposition of thrust sheet units made up from the base to the top (Fig. 6) by: Neritic unit (Cretaceous

carbonate), Ultra-Tellian unit (Cretaceous-Eocene marls and marly limestone), Tellian s.s. (sensu-structo): Marly dominance (Cretaceous-Eocene), Numidian unit with sandstone Burdigalian, clay and flysch (Eocene), Mio-Plio-Quaternary: sandy clays, marls and conglomerate (Mio-Pliocene). Alluvial terraces and lacustrine calcareous formations with Quaternary age (Raoult, 1974; Coiffait and Vila, 1977; Coiffait, 1992; Vila, 1980). This edifice was deposited during paroxysmal compressional phases Eocene and Miocene (Durand-Delga, 1969; Raoult, 1974).




Fig. 5: Oro-hydrographic and toponymical map of the study area. (After BENZID Y, 2017 modified)




Fig. 6: Geological position of the area study (Q: Quaternary undifferentiated (Villafranchien to current); AS: Allochthonous South Sétifien group; D: Limestone chain: Paleozoic to Upper Lutetian of the internal, median and external zones; FM: Mauritanian flysches; Fm:

Massylian flyschs, Typical series with phanites or green and red breccias; G: Granite, granodiorite, microgranite, dolerites and rhyolites; N: Constantine neritic tablecloth; NT: S.S tellian tablecloths; Nt: Tellian tablecloths; OMK: Kabyle oligo- Miocene, olistostrome; P: Marine Pliocene of coastal areas; PT: Penile tablecloth; S: Allochthonous Sellaoua-type training; SB: Basic base of Cap Bougaroun (peridotites and retromorphosed kinzigites); SE: Pedestal of Jebel Edough; SK: Metamorphic base of Little Kabylia and Bou Hatem (phyllades, mica schists, porphyroids, gneiss); SS: Allochthonous and parautochthonous foreland formations; TB: Effusive rocks; gN: Numidian sandstone and mixed series; m: Undifferentiated transgressive marine Miocene of the tellian aquifers and of the allochthonous, parautochthonous foreland; m': Upper Burdigalien –Langhien of the littoral margin; mp: Continental Mio- Pliocene; t: Triassic (Clays, ground gypsum and calcareo-dolomitic ice cubes); u: Ultra-telliene tablecloth - Typical series (Neocomian to Upper Lute-tian) marly and marl-limestone to light Cretaceous)




Fig. 7: Situation of the Safsaf watershed




Fig. 8: Situation of the Rhumel Smendou watershed




Fig. 9: Localisation of the Rhumel-Smendou and Safsaf watershed and regressive erosion, A - Rhumel Smendou watershed, B - Safsaf watershed

Methodology of producing the fluvial geomorphology map in the North-East of Algeria – identification of vulnerability factors

The elaboration of the fluvial-geomorphology map of the North East of Algeria was carried out based on:

- The map of the hydro-climatological network made by the national agency of hydraulic resources (ANRH) on a scale of 1/500000 (2006 edition).
- The "Shutter Radar Topography Mission" SRTM was used to delineate a digital elevation model (DEM) with a resolution of 30 meters.

In order to process this digital elevation model, ArcGIS 10.8 software was used to generate the

watershed, sub-watershed and hydrography network of the North-East of Algeria which are the Constantine coastal watershed (Fig. 7, 8, 10), the Kebir Rhumel watershed, and the Sybouse watershed, the studied area is divided into three Zones:

- The first zone is the Constantine coastal zone located on the northern slope of the Numidic chain (Fig. 7, 10).
- The second zone is the intramountain zone (Fig. 9, 10); situated between the coastal and the north slope of Numidic chain zone.
- The third zone is the interior zone limited between the south slope of Numidic chain from the north and Constantine mountains to the south (Fig. 8, 10).




Fig. 10: The fluvial geomorphology map of the North East of Algeria

This important map shows the genesis and the evolution of the hydrographic network of the Maghribides chain (North-East of Algeria). There is a concordance between the mountains and the hydrographic network, the rivers in longitudinal with the mountains (Numidic chain) take the East-West direction. Moreover, the transversal rivers have a South-North direction.

It is necessary to take into consideration the heritage and the geological history during the evolution of the relief. The active dynamic of northern Algeria and the eastern part in particular, favours the genesis and the evolution of instabilities, which are mostly caused by terrain deformations. A complex morpho-structural context frequently promotes and

guides these deformations (Benabbas, 2006, Benzid, 2017; Benabbas et al., 2018).

The watersheds of the North-East of Algeria form a Mediterranean region with a large variety of erosional processes. The Pliocene topography formed the basis for the evolution of the hydrographic network of the North-East of Algerian Watersheds. After that, tectonic activities gave the region its mountainous appearance (Maghribides chain) which develops progressively at the end of the Pliocene. These active tectonic movements are also responsible for the resumption of river down cutting, which may incise deep valleys and gorges. (Khanchoul and Saaidia, 2017).



Fig. 11: A - Coastal zone in Collo; B - Intramountain zone

The role of the slope in the dynamics of the river

Energy component of water masses (dependence on velocity, gravity, depth of current and absolute absence) (kinetic energy is equal to $1/2 (mv^2)$).

1. Various longitudinal profile on the slope depending on the relief shape (Fig. 12 A, B);

- 2. River level;
- 3. River processes - erosion and accumulation (Fig. 4);
- 4. Analysis of deposits - reliefs (reeds, islets) and Granulometry (sampling, laboratory analysis);
- 5. The islets of the Buzău River in the Întorsura valley.




Fig. 12: A - Longitudinal profile: Buzău; B - Longitudinal profile: Rhumel

Tectonic risk within the Curvature Carpathians

The Buzău River basin is located in a region known for its tectonic mobility specific to the Curvature Carpathians (Visarion et al., 1977). This tectonic mobility is maintained by the frequency of earthquakes in the Vrancea seismic region, with earthquakes of more

than 7 degrees Richter. Each earthquake is a risk factor for the triggering of geomorphological processes of the watershed and the riverbed.

The mountainous and Sub-Carpathian region is part of the sedimentary flysch of the Cretaceous, Paleogene, Mio-Pliocene and plain unit. The Quaternary with fluvio-lacustrine deposits, is dominated by